## **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:
C12N 15/86, 15/35, 5/10, A61K 48/00
A2
(11) International Publication Number: WO 00/28061
(43) International Publication Date: 18 May 2000 (18.05.00)

US

(21) International Application Number: PCT/US99/25694
 (22) International Filing Date: 2 November 1999 (02.11.99)

(30) Priority Data:

(71) Applicant (for all designated States except US): THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104–3147 (US).

5 November 1998 (05.11.98)

(72) Inventors; and

60/107,114

(75) Inventors/Applicants (for US only): WILSON, James, M. [US/US]; 1350 N. Avignon Drive, Gladwyne, PA 19035 (US). XIAO, Weidong [CN/US]; Apartment P4, 155 Washington Lane, Jenkintown, PA 19046 (US).

(74) Agents: KODROFF, Cathy, A. et al.; Howson & Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

(54) Title: ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

#### (57) Abstract

The nucleic acid sequences of adeno-associated virus (AAV) serotype 1 are provided, as are vectors and host cells containing these sequences and functional fragments thereof. Also provided are methods of delivering genes via AAV-1 derived vectors.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

			•				
AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
ΛM	Armenia	Fi	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	Prance	LŲ	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ.	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary ·	ML	Mali	TT	Trinidad and Tobago
BJ	Benin '	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Салада	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF ·	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Келуа	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan -	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	244	23/1102DWC
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		•
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 00/28061 PCT/US99/25694

# ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

This work was supported by the National Institutes of Health, grant no. P30 DK47757-06 and PO1 HD32649-04. The US government may have certain rights in this invention.

#### Field of the Invention

5

This invention relates generally to viral vector, and more particularly, to recombinant viral vectors useful for gene delivery.

#### Background of the Invention

10 Adeno-associated viruses are small, single-stranded DNA viruses which require helper virus to facilitate efficient replication [K.I. Berns, Parvoviridae: the viruses and their replication, p. 1007-1041, in F.N. Fields et al., Fundamental virology, 3rd ed., vol. 2, (Lippencott-Raven Publishers, Philadelphia, PA) (1995)]. The 4.7 kb genome of AAV is characterized by two inverted terminal repeats (ITR) and two open reading frames which encode the Rep proteins and Cap proteins, 15 respectively. The Rep reading frame encodes four proteins of molecular weight 78 kD, 68 kD, 52 kD and 40 kD. These proteins function mainly in regulating AAV replication and integration of the AAV into a host cell's chromosomes. The Cap reading frame encodes three structural proteins in molecular weight 85 kD (VP 1), 72 kD (VP2) and 61 kD (VP3) [Berns, cited above]. More than 80% of total proteins in 20 AAV virion comprise VP3. The two ITRs are the only cis elements essential for AAV replication, packaging and integration. There are two conformations of AAV ITRs called "flip" and "flop". These differences in conformation originated from the replication model of adeno-associated virus which use the ITR to initiate and reinitiate 25 the replication [R.O. Snyder et al., J. Virol., 67:6096-6104 (1993); K.I. Berns. Microbiological Reviews, 54:316-329 (1990)].

AAVs have been found in many animal species, including primates, canine, fowl and human [F.A. Murphy et al., "The Classification and Nomenclature of Viruses: Sixth Report of the International Committee on Taxonomy of Viruses",

10

25

Archives of Virology, (Springer-Verlag, Vienna) (1995)]. In addition to five known primate AAVs (AAV-1 to AAV-5), AAV-6, another serotype closely related to AAV-2 and AAV-1 has also been isolated [E. A. Rutledge et al., J. Virol., 72:309-319 (1998)]. Among all known AAV serotypes, AAV-2 is perhaps the most well-characterized serotype, because its infectious clone was the first made [R.J. Samulski et al., Proc. Natl., Acad. Sci. USA, 79:2077-2081 (1982)]. Subsequently, the full sequences for AAV-3A, AAV-3B, AAV-4 and AAV-6 have also been determined [Rutledge, cited above; J.A.Chiorini et al., J. Virol., 71:6823-6833 (1997); S. Muramatsu et al., Virol., 221:208-217 (1996)]. Generally, all AAVs share more than 80% homology in nucleotide sequence.

A number of unique properties make AAV a promising vector for human gene therapy [Muzyczka, Current Topics in Microbiology and Immunology, 158:97-129 (1992)]. Unlike other viral vectors, AAVs have not been shown to be associated with any known human disease and are generally not considered pathogenic. Wild type

15 AAV is capable of integrating into host chromosomes in a site specific manner [R. M. Kotin et al., Proc, Natl. Acad, Sci, USA, 87:2211-2215 (1990)- R.J. Samulski, EMBO J., 10(12):3941-3950 (1991)]. Recombinant AAV vectors can integrate into tissue cultured cells in chromosome 19 if the rep proteins are supplied in *trans* [C. Balague et al., J. Virol., 71:3299-3306 (1997); R. T. Surosky et al., J. Virol., 71:7951-7959 (1997)]. The integrated genomes of AAV have been shown to allow long term gene expression in a number of tissues, including, muscle, liver, and brain [K. J. Fisher, Nature Med., 3(3):306-312 (1997); R. 0. Snyder et al., Nature Genetics, 16:270-276 (1997), X. Xiao et al., Experimental Neurology, 144:113-124

AAV-2 has been shown to be present in about 80-90% of the human population. Earlier studies showed that neutralizing antibodies for AAV-2 are prevalent [W. P. Parks et al., <u>J. Virol.</u>, <u>2</u>:716-722 (1970)]. The presence of such antibodies may significantly decrease the usefulness of AAV vectors based on AAV-2 despite its other merits. What are needed in the art are vectors characterized by the

(1997); Xiao, J. Virol., 70(11):8098-8108 (1996)].

10

15

20

25

advantages of AAV-2, including those described above, without the disadvantages, including the presence of neutralizing antibodies.

#### Summary of the Invention

In one aspect, the invention provides an isolated AAV-1 nucleic acid molecule which is selected from among SEQ ID NO: 1, the strand complementary to SEQ ID NO: 1, and cDNA and RNA sequences complementary to SEQ ID NO: 1 and its complementary strand.

In another aspect, the present invention provides AAV ITR sequences, which include the 5' ITR sequences, nt 1 to 143 of SEQ ID NO: 1; the 3' ITR sequences, nt 4576 to 4718 of SEQ ID NO: 1, and fragments thereof.

In yet another aspect, the present invention provides a recombinant vector comprising an AAV-1 ITR and a selected transgene. Preferably, the vector comprises both the 5' and 3' AAV-1 ITRs between which the selected transgene is located.

In still another aspect, the invention provides a recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.

In a further aspect, the present invention provides a nucleic acid molecule encoding an AAV-1 rep coding region and an AAV-1 cap coding region.

In still another aspect, the present invention provides a host cell transduced with a recombinant viral vector of the invention. The invention further provides a host cell stably transduced with an AAV-1 P5 promoter of the invention.

In still a further aspect, the present invention provides a pharmaceutical composition comprising a carrier and a vector of the invention.

In yet another aspect, the present invention provides a method for AAV-mediated delivery of a transgene to a host involving the step of delivering to a selected
host a recombinant viral vector comprising a selected transgene under the control of
sequences which direct expression thereof and an adeno-associated virus 1 (AAV-1)
virion.

In another aspect, the invention provides a method for in vitro production of a selected gene product using a vector of the invention.

Other aspects and advantages of the invention will be readily apparent to one of skill in the art from the detailed description of the invention.

# 5 Brief Description of the Drawings

10

15

20

25

Figs. 1A-1C illustrate the alignment of nucleotides of AAV-1 [SEQ ID NO: 1], AAV-2 [SEQ ID NO: 18] and AAV-6 [SEQ ID NO: 19]. The alignment was done with MacVector 6.0. The full sequences of AAV-1 are shown in the top line. Nucleotides in AAV-2 and AAV-6 identical to AAV-1 are symbolized by "." and gaps by "-". Some of the conserved features among AAVs are marked in this figure. Note the 3' ITRs of AAV-1 and AAV-6 are shown in different orientations.

Fig. 2 illustrates the predicted secondary structure of AAV-1 ITR. The nucleotides in AAV-2 and AAV-6 are shown in italic and bold respectively.

Fig. 3A illustrates a hypothesis of how AAV-6 arose from the homologous recombination between AAV-1 and AAV-2. The major elements of AAV-1 are indicated in the graph. A region that is shared between AAV-1, AAV-2 and AAV-6 is shown in box with waved lines.

Fig. 3B is a detailed illustration of a 71 bp homologous region among AAV-1, AAV-2 and AAV-6. Nucleotides that differ among these serotypes are indicated by arrows.

Fig. 4A is a bar chart illustrating expression levels of human alpha 1 antitrypsin (α1AT) in serum following delivery of hAAT via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 4B is a bar chart illustrating expression levels of erythropoietin (epo) in serum following delivery of the epo gene via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 5A is a bar chart illustrating expression levels of  $\alpha 1$  AT in liver following delivery of  $\alpha 1$  AT as described in Example 7.

20

25

Fig. 5B is a bar chart demonstrating expression levels of epo in liver following delivery of epo as described in Example 7.

Fig. 5C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to liver as described in Example 7.

Fig. 5D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of a1AT or epo to liver as described in Example 7.

Fig. 6A is a bar chart illustrating expression levels of  $\alpha 1$  AT in muscle following delivery of  $\alpha 1$  AT as described in Example 7.

Fig. 6B is a bar chart demonstrating expression levels of epo in muscle following delivery of epo as described in Example 7.

Fig. 6C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to muscle as described in Example 7.

Fig. 6D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of a1AT or epo to muscle as described in Example 7.

## 15 <u>Detailed Description of the Invention</u>

The present invention provides novel nucleic acid sequences for an adeno-associated virus of serotype 1 (AAV-1). Also provided are fragments of these AAV-1 sequences. Among particularly desirable AAV-1 fragments are the inverted terminal repeat sequences (ITRs), rep and cap. Each of these fragments may be readily utilized, e.g., as a cassette, in a variety of vector systems and host cells. Such fragments may be used alone, in combination with other AAV-1 sequences or fragments, or in combination with elements from other AAV or non-AAV viral sequences. In one particularly desirable embodiment, a cassette may contain the AAV-1 ITRs of the invention flanking a selected transgene. In another desirable embodiment, a cassette may contain the AAV-1 rep and/or cap proteins, e.g., for use in producing recombinant (rAAV) virus.

Thus, the AAV-1 sequences and fragments thereof are useful in production of rAAV, and are also useful as antisense delivery vectors, gene therapy vectors, or vaccine vectors. The invention further provides nucleic acid molecules, gene delivery

10

15

20

25

vectors, and host cells which contain the AAV-1 sequences of the invention. Also provided a novel methods of gene delivery using AAV vectors.

As described herein, the vectors of the invention containing the AAV-1 capsid proteins of the invention are particularly well suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other AAV serotype based vectors, as well as other viral vectors. The rAAV vectors of the invention are particularly advantageous in rAAV readministration and repeat gene therapy.

These and other embodiments and advantages of the invention are described in more detail below. As used throughout this specification and the claims, the term "comprising" is inclusive of other components, elements, integers, steps and the like.

# I. AAV-1 NUCLEIC ACID AND PROTEIN SEQUENCES

The AAV-1 nucleic acid sequences of the invention include the DNA sequences of SEQ ID NO: 1 (Figs. 1A-1C), which consists of 4718 nucleotides. The AAV-1 nucleic acid sequences of the invention further encompass the strand which is complementary to SEQ ID NO: 1, as well as the RNA and cDNA sequences corresponding to SEQ ID NO: 1 and its complementary strand. Also included in the nucleic acid sequences of the invention are natural variants and engineered modifications of SEQ ID NO: 1 and its complementary strand. Such modifications include, for example, labels which are known in the art, methylation, and substitution of one or more of the naturally occurring nucleotides with an analog.

Further included in this invention are nucleic acid sequences which are greater than 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 98 - 99% identical or homologous to SEQ ID NO:1. The term "percent sequence identity" or "identical" in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length sequence, or a fragment at least about nine nucleotides, usually at least about 20 - 24 nucleotides, at least about 28 - 32 nucleotides, and preferably at least about 36 or more nucleotides. There are a number of different

10

15

20

25

30

algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using Fasta, a program in GCG Version 6.1. Fasta provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, 1990, herein incorporated by reference). For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference.

The term "substantial homology" or "substantial similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 - 99% of the sequence.

Also included within the invention are fragments of SEO ID NO: 1, its complementary strand, cDNA and RNA complementary thereto. Suitable fragments are at least 15 nucleotides in length, and encompass functional fragments which are of biological interest. Certain of these fragments may be identified by reference to Figs. 1A-1C. Examples of particularly desirable functional fragments include the AAV-1 inverted terminal repeat (ITR) sequences of the invention. In contrast to the 145 nt ITRs of AAV-2, AAV-3, and AAV-4, the AAV-1 ITRs have been found to consist of only 143 nucleotides, yet advantageously are characterized by the T-shaped hairpin structure which is believed to be responsible for the ability of the AAV-2 ITRs to direct site-specific integration. In addition, AAV-1 is unique among other AAV serotypes, in that the 5' and 3' ITRs are identical. The full-length 5' ITR sequences of AAV-1 are provided at nucleotides I-143 of SEQ ID NO: 1 (Fig. 1A) and the fulllength 3' ITR sequences of AAV-1 are provided at nt 4576-4718 of SEQ ID NO: 1 (Fig. 1C). One of skill in the art can readily utilize less than the full-length 5' and/or 3' 1TR sequences for various purposes and may construct modified ITRs using conventional techniques, e.g., as described for AAV-2 ITRs in Samulski et al, Cell, 33:135-143 (1983).

10

15

20

25

Another desirable functional fragment of the AAV-1 genome is the P5 promoter of AAV-1 which has sequences unique among AAV P5 promoters, while maintaining critical regulatory elements and functions. This promoter is located within nt 236 - 299 of SEQ ID NO: 1 (Fig. 1A). Other examples of functional fragments of interest include the sequences at the junction of the rep/cap, e.g., the sequences spanning nt 2306-2223, as well as larger fragments which encompass this junction which may comprise 50 nucleotides on either side of this junction. Still other examples of functional fragments include the sequences encoding the rep proteins. Rep 78 is located in the region of nt 334 - 2306 of SEQ ID NO: 1; Rep 68 is located in the region of nt 334-2272, and contains an intron spanning nt 1924-2220 of SEQ ID NO: 1. Rep 52 is located in the region of nt 1007 - 2304 of SEQ ID NO: 1; rep 40 is located in the region of nt 1007 - 2272, and contains an intron spanning nt 1924-2246 of SEQ ID NO: 1. Also of interest are the sequences encoding the capsid proteins, VP 1 [nt 2223-4431 of SEQ ID NO: 1], VP2 [nt 2634-4432 of SEQ ID NO: 1] and VP3 [nt 2829-4432 of SEQ ID NO: 1]. Other fragments of interest may include the AAV-1 P19 sequences, AAV-1 P40 sequences, the rep binding site, and the terminal resolute site (TRS).

The invention further provides the proteins and fragments thereof which are encoded by the AAV-1 nucleic acids of the invention. Particularly desirable proteins include the rep and cap proteins, which are encoded by the nucleotide sequences identified above. These proteins include rep 78 [SEQ ID NO:5], rep 68 [SEQ ID NO:7], rep 52 [SEQ ID NO:9], rep 40 [SEQ ID NO: 11], vpl [SEQ ID NO: 13], vp2 [SEQ ID NO: 15], and vp3 [SEQ IID NO: 17] and functional fragments thereof while the sequences of the rep and cap proteins have been found to be closely related to those of AAV-6, there are differences in the amino acid sequences (see Table 1 below), as well as differences in the recognition of these proteins by the immune system. However, one of skill in the art may readily select other suitable proteins or protein fragments of biological interest. Suitably, such fragments are at least 8 amino acids in length. However, fragments of other desired lengths may be readily utilized.

Such fragments may be produced recombinantly or by other suitable means, e.g., chemical synthesis.

The sequences, proteins, and fragments of the invention may be produced by any suitable means, including recombinant production, chemical synthesis, or other synthetic means. Such production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

#### II. VIRAL VECTORS

5

10

15

20

25

In another aspect, the present invention provides vectors which utilize the AAV-1 sequences of the invention, including fragments thereof, for delivery of a heterologous gene or other nucleic acid sequences to a target cell. Suitably, these heterologous sequences (i.e., a transgene) encode a protein or gene product which is capable of being expressed in the target cell. Such a transgene may be constructed in the form of a "minigene". Such a "minigene" includes selected heterologous gene sequences and the other regulatory elements necessary to transcribe the gene and express the gene product in a host cell. Thus, the gene sequences are operatively linked to regulatory components in a manner which permit their transcription. Such components include conventional regulatory elements necessary to drive expression of the transgene in a cell containing the viral vector. The minigene may also contain a selected promoter which is linked to the transgene and located, with other regulatory elements, within the selected viral sequences of the recombinant vector.

Selection of the promoter is a routine matter and is not a limitation of this invention. Useful promoters may be constitutive promoters or regulated (inducible) promoters, which will enable control of the timing and amount of the transgene to be expressed. For example, desirable promoters include the cytomegalovirus (CMV) immediate early promoter/enhancer [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the Rous sarcoma virus LTR promoter/enhancer, and the chicken cytoplasmic β-actin promoter [T. A. Kost et al, Nucl, Acids Res., 11(23):8287 (1983)]. Still other desirable promoters are the albumin promoter and an AAV P5 promoter. Optionally, the selected promoter is used in conjunction with a heterologous enhancer, e.g., the β-

WO 00/28061 PCT/US99/25694

10

actin promoter may be used in conjunction with the CMV enhancer. Yet other suitable or desirable promoters and enhancers may be selected by one of skill in the art.

5

10

15

20

25

30

The minigene may also desirably contain nucleic acid sequences heterologous to the viral vector sequences including sequences providing signals required for efficient polyadenylation of the transcript (poly-A or pA) and introns with functional splice donor and acceptor sites. A common poly-A sequence which is employed in the exemplary vectors of this invention is that derived from the papovavirus SV-40. The poly-A sequence generally is inserted in the minigene downstream of the transgene sequences and upstream of the viral vector sequences. A common intron sequence is also derived from SV-40, and is referred to as the SV40 T intron sequence. A minigene of the present invention may also contain such an intron, desirably located between the promoter/enhancer sequence and the transgene. Selection of these and other common vector elements are conventional [see, e.g., Sambrook et al, "Molecular Cloning. A Laboratory Manual", 2d edit., Cold Spring Harbor Laboratory, New York (1989) and references cited therein] and many such sequences are available from commercial and industrial sources as well as from Genebank.

The selection of the transgene is not a limitation of the present invention. Suitable transgenes may be readily selected from among desirable reporter genes, therapeutic genes, and optionally, genes encoding immunogenic polypeptides. Examples of suitable reporter genes include  $\beta$ -galactosidase ( $\beta$ -gal), an alkaline phosphatase gene, and green fluorescent protein (GFP). Examples of therapeutic genes include, cytokines, growth factors, hormones, and differentiation factors, among others. The transgene may be readily selected by one of skill in the art. See, e.g., WO 98/09657, which identifies other suitable transgenes.

Suitably, the vectors of the invention contain, at a minimum, cassettes which consist of fragments of the AAV-1 sequences and proteins. In one embodiment, a vector of the invention comprises a selected transgene, which is flanked by a 5' ITR and a 3' ITR, at least one of which is an AAV-1 ITR of the invention. Suitably,

vectors of the invention may contain a AAV-1 P5 promoter of the invention. In yet another embodiment, a plasmid or vector of the invention contains AAV-1 rep sequences. In still another embodiment, a plasmid or vector of the invention contains at least one of the AAV-1 cap proteins of the invention. Most suitably, these AAV-1-derived vectors are assembled into viral vectors, as described herein.

## A. AAV Viral Vectors

5

10

15

20

25

In one aspect, the present invention provides a recombinant AAV-1 viral vector produced using the AAV-1 capsid proteins of the invention. The packaged rAAV-1 virions of the invention may contain, in addition to a selected minigene, other AAV-1 sequences, or may contain sequences from other AAV serotypes.

Methods of generating rAAV virions are well known and the selection of a suitable method is not a limitation on the present invention. See, e.g., K. Fisher et al, <u>J. Virol.</u>, <u>70</u>:520-532 (1993) and US Patent 5,478,745. In one suitable method, a selected host cell is provided with the AAV sequence encoding a rep protein, the gene encoding the AAV cap protein and with the sequences for packaging and subsequent delivery. Desirably, the method utilizes the sequences encoding the AAV-1 rep and/or cap proteins of the invention.

In one embodiment, the rep/cap genes and the sequences for delivery are supplied by co-transfection of vectors carrying these genes and sequences. In one currently preferred embodiment, a cis (vector) plasmid, a trans plasmid containing the rep and cap genes, and a plasmid containing the adenovirus helper genes are co-transfected into a suitable cell line, e.g., 293. Alternatively, one or more of these functions may be provided in trans via separate vectors, or may be found in a suitably engineered packaging cell line.

An exemplary cis plasmid will contain, in 5' to 3' order, AAV 5' ITR, the selected transgene, and AAV 3' ITR. In one desirable embodiment, at least one of the AAV ITRs is a 143 nt AAV-1 ITR. However, other AAV serotype ITRs may be readily selected. Suitably, the full-length ITRs are utilized. However, one of skill in

WO 00/28061 PCT/US99/25694

12

the art can readily prepare modified AAV ITRs using conventional techniques. Similarly, methods for construction of such plasmids is well known to those of skill in the art.

A trans plasmid for use in the production of the rAAV-1 virion particle may be prepared according to known techniques. In one desired embodiment, this plasmid contains the rep and cap proteins of AAV-1, or functional fragments thereof. Alternatively, the rep sequences may be from another selected AAV serotype.

5

10

15

20

25

30

The cis and trans plasmid may then be co-transfected with a wild-type helper virus (e.g., Ad2, Ad5, or a herpesvirus), or more desirably, a replication - defective adenovirus, into a selected host cell. Alternatively, the cis and trans plasmid may be co-transfected into a selected host cell together with a transfected plasmid which provides the necessary helper functions. Selection of a suitable host cell is well within the skill of those in the art and include such mammalian cells as 293 cells, HeLa cells, among others.

Alternatively, the cis plasmid and, optionally the trans plasmid, may be transfected into a packaging cell line which provides the remaining helper functions necessary for production of a rAAV containing the desired AAV-1 sequences of the invention. An example of a suitable packaging cell line, where an AAV-2 capsid is desired, is B-50, which stably expresses AAV-2 rep and cap genes under the control of a homologous P5 promoter. This cell line is characterized by integration into the cellular chromosome of multiple copies (at least 5 copies) of P5-rep-cap gene cassettes in a concatomer form. This B-50 cell line was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, on September 18, 1997 under Accession No. CRL-12401 pursuant to the provisions of the Budapest Treaty. However, the present invention is not limited as to the selection of the packaging cell line.

Exemplary transducing vectors based on AAV-1 capsid proteins have been tested both *in vivo and in vitro*, as described in more detail in Example 4. In these studies, it was demonstrated that recombinant AAV vector with an AAV-1 virion can transduce both mouse liver and muscle. These, and other AAV-1 based

WO 00/28061 PCT/US99/25694

13

gene therapy vectors which may be generated by one of skill in the art are beneficial for gene delivery to selected host cells and gene therapy patients since the neutralization antibodies of AAV-1 present in much of the human population exhibit different patterns from other AAV serotypes and therefore do not neutralize the AAV-1 virions. One of skill in the art may readily prepare other rAAV viral vectors containing the AAV-1 capsid proteins provided herein using a variety of techniques known to those of skill in the art. One may similarly prepare still other rAAV viral vectors containing AAV-1 sequence and AAV capsids of another serotype.

#### B. Other Viral Vectors

the AAV-1 cap, and the AAV-1 P5 promoter sequences.

the present invention is not limited to any particular method.

5

15

20

25

30

One of skill in the art will readily understand that the AAV-1 sequences of the invention can be readily adapted for use in these and other viral vector systems for in vitro, ex vivo or in vivo gene delivery. Particularly well suited for use in such viral vector systems are the AAV-1 ITR sequences, the AAV-1 rep,

For example, in one desirable embodiment, the AAV-1 ITR sequences of the invention may be used in an expression cassette which includes AAV-1 5' ITR, a non-AAV DNA sequences of interest (e.g., a minigene), and 3' ITR and which lacks functional rep/cap. Such a cassette containing an AAV-1 ITR may be located on a plasmid for subsequent transfection into a desired host cell, such as the cis plasmid described above. This expression cassette may further be provided with an AAV capsid of a selected serotype to permit infection of a cell or stably transfected into a desired host cell for packaging of rAAV virions. Such an expression cassette may be readily adapted for use in other viral systems, including adenovirus systems and lentivirus systems. Methods of producing Ad/AAV vectors are well known to those of skill in the art. One desirable method is described in PCT/US95/14018. However,

Another aspect of the present invention is the novel AAV-1 P5 promoter sequences which are located in the region spanning at 236 - 299 of SEQ ID NO: 1. This promoter is useful in a variety of viral vectors for driving expression of a desired transgene.

10

15

20

25

Similarly, one of skill in the art can readily select other fragments of the AAV-1 genome of the invention for use in a variety of vector systems. Such vectors systems may include, e.g., lentiviruses, retroviruses, poxviruses, vaccinia viruses, and adenoviral systems, among others. Selection of these vector systems is not a limitation of the present invention.

# C. Host Cells And Packaging Cell Lines

In yet another aspect, the present invention provides host cells which may be transiently transfected with AAV-1 nucleic acid sequences of the invention to permit expression of a desired transgene or production of a rAAV particle. For example, a selected host cell may be transfected with the AAV-1 P5 promoter sequences and/or the AAV-1 5' ITR sequences using conventional techniques. Providing AAV helper functions to the transfected cell lines of the invention results in packaging of the rAAV as infectious rAAV particles. Such cell lines may be produced in accordance with known techniques [see, e.g, US Patent No. 5,658,785], making use of the AAV-1 sequences of the invention.

Alternatively, host cells of the invention may be stably transfected with a rAAV expression cassette of the invention, and with copies of AAV-1 rep and cap genes. Suitable parental cell lines include mammalian cell lines and it may be desirable to select host cells from among non-simian mammalian cells. Examples of suitable parental cell lines include, without limitation, HeLa [ATCC CCL 2], A549 [ATCC Accession No. CCL 185], KB [CCL 17], Detroit [e.g., Detroit 510, CCL 72] and WI-38 [CCL 75] cells. These cell lines are all available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 USA. Other suitable parent cell lines may be obtained from other sources and may be used to construct stable cell lines containing the P5 and/or AAV rep and cap sequences of the invention.

Recombinant vectors generated as described above are useful for delivery of the DNA of interest to cells.

10

15

20

25

#### III. METHODS OF DELIVERING GENES VIA AAV-1 DERIVED VECTORS

In another aspect, the present invention provides a method for delivery of a transgene to a host which involves transfecting or infecting a selected host cell with a recombinant viral vector generated with the AAV-1 sequences (or functional fragments thereof) of the invention. Methods for delivery are well known to those of skill in the art and are not a limitation of the present invention.

In one desirable embodiment, the invention provides a method for AAV-mediated delivery of a transgene to a host. This method involves transfecting or
infecting a selected host cell with a recombinant viral vector containing a selected
transgene under the control of sequences which direct expression thereof and AAV-1
capsid proteins.

Optionally, a sample from the host may be first assayed for the presence of antibodies to a selected AAV serotype. A variety of assay formats for detecting neutralizing antibodies are well known to those of skill in the art. The selection of such an assay is not a limitation of the present invention. See, e.g., Fisher et al, Nature Med., 3(3):306-312 (March 1997) and W. C. Manning et al, Human Gene Therapy, 9:477-485 (March 1, 1998). The results of this assay may be used to determine which AAV vector containing capsid proteins of a particular serotype are preferred for delivery, e.g., by the absence of neutralizing antibodies specific for that capsid serotype.

In one aspect of this method, the delivery of vector with AAV-1 capsid proteins may precede or follow delivery of a gene via a vector with a different serotype AAV capsid protein. Thus, gene delivery via rAAV vectors may be used for repeat gene delivery to a selected host cell. Desirably, subsequently administered rAAV vectors carry the same transgene as the first rAAV vector, but the subsequently administered vectors contain capsid proteins of serotypes which differ from the first vector. For example, if a first vector has AAV-2 capsid proteins, subsequently administered vectors may have capsid proteins selected from among the other serotypes, including AAV-1, AAV-3A, AAV-3B, AAV-4 and AAV-6.

WO 00/28061

5

10

15

20

25

30

Thus, a rAAV-1-derived recombinant viral vector of the invention provides an efficient gene transfer vehicle which can deliver a selected transgene to a selected host cell in vivo or ex vivo even where the organism has neutralizing antibodies to one or more AAV serotypes. These compositions are particularly well suited to gene delivery for therapeutic purposes. However, the compositions of the invention may also be useful in immunization. Further, the compositions of the invention may also be used for production of a desired gene product in vitro.

The above-described recombinant vectors may be delivered to host cells according to published methods. An AAV viral vector bearing the selected transgene may be administered to a patient, preferably suspended in a biologically compatible solution or pharmaceutically acceptable delivery vehicle. A suitable vehicle includes sterile saline. Other aqueous and non-aqueous isotonic sterile injection solutions and aqueous and non-aqueous sterile suspensions known to be pharmaceutically acceptable carriers and well known to those of skill in the art may be employed for this purpose.

The viral vectors are administered in sufficient amounts to transfect the cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse effects, or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the liver, oral, intranasal, intravenous, intramuscular, subcutaneous, intradermal, and other parental routes of administration. Routes of administration may be combined, if desired.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective human dosage of the viral vector is generally in the range of from about 1 ml to about 100 ml of solution containing concentrations of from about 1 x 10<sup>9</sup> to 1 x 10<sup>16</sup> genomes virus vector. A preferred human dosage may be about 1 x 10<sup>13</sup> to 1 x 10<sup>16</sup> AAV genomes. The dosage will be adjusted to balance the therapeutic benefit against any side effects and

10

15

20

25

such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed. The levels of expression of the transgene can be monitored to determine the frequency of dosage resulting in viral vectors, preferably AAV vectors containing the minigene. Optionally, dosage regimens similar to those described for therapeutic purposes may be utilized for immunization using the compositions of the invention. For *in vitro* production, a desired protein may be obtained from a desired culture following transfection of host cells with a rAAV containing the gene encoding the desired protein and culturing the cell culture under conditions which permits expression. The expressed protein may then be purified and isolated, as desired. Suitable techniques for transfection, cell culturing, purification, and isolation are known to those of skill in the art.

The following examples illustrate several aspects and embodiments of the invention.

#### Example 1 - Generation of Infectious Clone of AAV-1

The replicated form DNA of AAV-1 was extracted from 293 cells that were infected by AAV-1 and wild type adenovirus type 5.

#### A. Cell Culture and Virus

AAV-free 293 cells and 84-31 cells were provided by the human application laboratory of the University of Pennsylvania. These cells were cultured in Dulbecco's Modified Eagle Medium with 10% fetal bovine serum (Hyclone), penicillin (100 U/ml) and streptomycin at 37°C in a moisturized environment supplied with 5% C0<sub>2</sub>. The 84-31 cell line constitutively expresses adenovirus genes E1a, Elb, E4/ORF6, and has been described previously [K. J. Fisher, J. Virol., 70:520-532 (1996)]. AAV-1 (ATCC VR-645) seed stock was purchased from American Type Culture Collection (ATCC, Manassas, VA). AAV viruses were propagated in 293 cells with wild type Ad5 as a helper virus.

#### B. Recombinant AAV Generation

The recombinant AAV viruses were generated by transfection using an adenovirus free method. Briefly, the cis plasmid (with AAV ITR), trans plasmid (with

WO 00/28061 PCT/US99/25694

18

AAV rep gene and cap gene) and helper plasmid (pFa13, with essential regions from the adenovirus genome) were simultaneously co-transfected into 293 cells in a ratio of 1:1:2 by calcium phosphate precipitation. The pFa13 helper plasmid has an 8 kb deletion in the adenovirus E2B region and has deletions in most of the late genes. This helper plasmid was generated by deleting the RsrII fragment from pFG140 (Microbix, Canada). Typically, 50 µg of DNA (cis:trans:PFa13 at ratios of 1:1:2, respectively) was transfected onto a 15 cm tissue culture dish. The cells were harvested 96 hours post-transfection, sonicated and treated with 0.5% sodium deoxycholate (37°C for 10 min). Cell lysates were then subjected to two rounds of a CsCl gradient. Peak fractions containing AAV vector were collected, pooled, and dialyzed against PBS before injecting into animals. To make rAAV virus with AAV-1 virion, the pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide rep and cap function.

5

10

15

20

25

30

For the generation of rAAV based on AAV-2, p5E18 was used as the trans plasmid since it greatly improved the rAAV yield. This plasmid, p5E18(2/2), expresses AAV-2 Rep and Cap and contains a P5 promoter relocated to a position 3' to the Cap gene, thereby minimizing expression of Rep78 and Rep68. The strategy was initially described by Li et al, J. Virol., 71:5236-5243 (1997). P5E18(2/2) was constructed in the following way. The previously described pMMTV-trans vector (i.e., the mouse mammary tumor virus promoter substituted for the P5 promoter in an AAV-2-based vector) was digested with SmaI and ClaI, filled in with the Klenow enzyme, and then recircularized with DNA ligase. The resulting construct was digested with XbaI, filled in, and ligated to the blunt-ended BamHI-XbaI fragment from pCR-p5, constructed in the following way. The P5 promoter of AAV was amplified by PCR and the amplified fragment was subsequently cloned into pCR2.1 (Invitrogen) to yield pCR-P5. The helper plasmid pAV1H was constructed by cloning the Bfal fragment of pAAV-2 into pBluescript II-SK(+) at the BcorV and Smal sites. The 3.0-kb XbaI-KpnI fragment from p5E18(2/2), the 2.3-kb XbaI-KpnI fragment from pAV1H, and the 1.7-kb KpnI fragment from pSE18(2/2) were incorporated into a separate plasmid P5E18(2/1), which contains AAV-2 Rep, AAV-1 Cap, and the

WO 00/28061 PCT/US99/25694

19

AAV-2 P5 promoter located 3' to the Cap gene. Plasmid p5E18(2/1) produced 10- to 20-fold higher quantities of the vector than pAV1H (i.e., 10<sup>12</sup> genomes/50 15-cm<sup>2</sup> plates).

## C. <u>DNA Techniques</u>

5

10

15

20

25

30

Hirt DNA extraction was performed as described in the art with minor modification [R.J. Samulski et al., Cell, 33:135-143 (1983)]. More particularly, Hirst solution without SDS was used instead of using original Hirt solution containing SDS. The amount of SDS present in the original Hirst solution was added after the cells had been fully suspended. To construct AAV-1 infectious clone, the Hirt DNA from AAV-1 infected 293 cells was repaired with Klenow enzyme (New England Biolabs) to ensure the ends were blunt. The treated AAV-1 Hirt DNA was then digested with BamHI and cloned into three vectors, respectively. The internal BamHI was cloned into pBlueScript II-SK+ cut with BamHI to get pAV1-BM. The left and right fragments were cloned into pBlueScript II-SK+ cut with BamHI + EcoRV to obtain pAV1-BL and pAV1-BR, respectively. The AAV sequence in these three plasmids were subsequently assembled into the same vector to get AAV-1 infectious clone pAAV-1. The helper plasmid for recombinant AAV-1 virus generation was constructed by cloning the Bfa I fragment of pAAV-1 into pBlueScript II-SK+ at the EcoRV site.

Analysis of the Hirt DNA revealed three bands, a dimer at 9.4 kb, a monomer at 4.7 kb and single-stranded DNA at 1.7 kb, which correlated to different replication forms of AAV-1. The monomer band was excised from the gel and then digested with *BamH*I. This resulted in three fragments of 1.1 kb, 0.8 kb and 2.8 kb. This pattern is in accordance with the description by Bantel-schaal and zur Hausen, Virol., 134(1):52-63 (1984). The 1.1 kb and 2.8 kb *BamH*I fragments were cloned into pBlueScript-KS(+) at *BamH*I and EcoRV site. The internal 0.8 kb fragment was cloned into *BamH*I site of pBlueScript-KS(+).

These three fragments were then subcloned into the same construct to obtain a plasmid (pAAV-1) that contained the full sequence of AAV-1. The pAAV-1 was then tested for its ability to rescue from the plasmid backbone and package

10

15

20

25

infectious virus. The pAAV-1 was then transfected to 293 cells and supplied with adenovirus type as helper at MOI 10. The virus supernatant was used to reinfect 293 cells.

For Southern blot analysis, Hirt DNA was digested with *DpnI* to remove bacteria-borne plasmid and probed with internal *BamHI* fragment of AAV-1. The membrane was then washed at high stringency conditions, which included: twice 30 minutes with 2X SSC, 0.1% SDS at 65°C and twice 30 minutes with 0.1X SSC, 0.1% SDS at 65°C. The membrane was then analyzed by both phosphor image and X-ray autoradiography. The results confirmed that pAAV-1 is indeed an infectious clone of AAV serotype 1.

#### Example 2 - Sequencing Analysis of AAV-1

The entire AAV-1 genome was then determined by automatic sequencing and was found to be 4718 nucleotides in length (Figs. 1A-1C). For sequencing, an ABI 373 automatic sequencer as used to determine the sequences for all plasmids and PCR fragments related to this study using the FS dye chemistry. All sequences were confirmed by sequencing both plus and minus strands. These sequences were also confirmed by sequencing two independent clones of pAV-BM, pAV-BL and pAV-BR. Since the replicated form of AAV-1 DNA served as the template for sequence determination, these sequences were also confirmed by sequencing a series of PCR products using original AAV-1 seed stock as a template.

The length of AAV-1 was found to be within the range of the other serotypes: AAV-3 (4726 nucleotides), AAV-4 (4774 nucleotides), AAV-2 (4681 nucleotides), and AAV-6 (4683 nucleotides).

The AAV-1 genome exhibited similarities to other serotypes of adenoassociated viruses. Overall, it shares more than 80% identity with other known AAV
viruses as determined by the computer program Megalign using default settings
[DNASTAR, Madison, WI]. The key features in AAV-2 can also be found in AAV1. First, AAV-1 has the same type of inverted terminal repeat which is capable of
forming T-shaped hairpin structures, despite the differences at the nucleotide level

WO 00/28061

5

10

15

20

25

30

(Figs. 2 and 3). The sequences of right ITRs and left ITRs of AAV-1 are identical. The AAV TR sequence is subdivided into A, A', B, B', C, C', D and D' [Bern, cited above].

These AAV ITR sequences are also virtually the same as those found in AAV-6 right ITR, there being one nucleotide difference in each of A and A' sequence, and the last nucleotide of the D sequence. Second, the AAV-2 rep binding motif [GCTCGCTCGCTCGCTG (SEQ ID NO: 20)] is well conserved. Such motif can also be found in the human chromosome 19 AAV-2 pre-integration region. Finally, non-structural and structural coding regions, and regulatory elements similar to those of other AAV serotypes also exist in AAV-1 genome.

Although the overall features of AAV terminal repeats are very much conserved, the total length of the AAV terminal repeat exhibits divergence. The terminal repeat of AAV-1 consists of 143 nucleotides while those of AAV-2, AAV-3, and AAV-4 are about 145 or 146 nucleotides. The loop region of AAV-1 ITR most closely resembles that of AAV-4 in that it also uses TCT instead of the TTT found in AAV-2 and AAV-3. The possibility of sequencing error was eliminated using restriction enzyme digestion, since these three nucleotides are part of the SacI site (gagete; nt 69-74 of SEQ ID NO: 1). The p5 promoter region of AAV-1 shows more variations in nucleotide sequences with other AAV serotypes. However, it still maintains the critical regulatory elements. The two copies of YY1 [See, Fig. 1A-1C] sites seemed to be preserved in all known AAV serotypes, which have been shown to be involved in regulating AAV gene expression. In AAV-4, there are 56 additional nucleotides inserted between YY1 and E-box/USF site, while in AAV-1, there are 26 additional nucleotides inserted before the E-box/USF site. The p19 promoter, p40 promoter and polyA can also be identified from the AAV-1 genome by analogy to known AAV serotypes, which are also highly conserved.

Thus, the analysis of AAV terminal repeats of various serotypes showed that the A and A' sequence is very much conserved. One of the reasons may be the Rep binding motif (GCTC)<sub>3</sub>GCTG [SEQ ID NO: 20]. These sequences appear to be essential for AAV DNA replication and site-specific integration. The same sequence

10

15

20

25

has also been shown to be preserved in a monkey genome [Samulski, personal communication]. The first 8 nucleotides of the D sequence are also identical in all known AAV serotypes. This is in accordance with the observation of the Srivastava group that only the first 10 nucleotides are essential for AAV packaging [X.S. Wang et al, J. Virol., 71:3077-3082 (1997); X.S. Wang et al, J. Virol., 71:1140-1146 (1997)]. The function of the rest of the D sequences still remain unclear. They may be somehow related to their tissue specificities. The variation of nucleotide in B and C sequence may also suggest that the secondary structure of the ITRs is more critical for its biological function, which has been demonstrated in many previous publications.

#### Example 3 - Comparison of AAV-1 Sequences

The nucleotide sequences of AAV-1, obtained as described above, were compared with known AAV sequences, including AAV-2, AAV-4 and AAV-6 using DNA Star Megalign. This comparison revealed a stretch of 71 identical nucleotides shared by AAV-1, AAV-2 and AAV-6. See, Figs. 1A-1C.

This comparison further suggested that AAV-6 is a hybrid formed by homologous recombination of AAV-1 and AAV-2. See, Figs. 3A and 3B. These nucleotides divide the AAV-6 genome into two regions. The 5' half of AAV-6 of 522 nucleotides is identical to that of AAV-2 except in 2 positions. The 3' half of AAV-6 including the majority of the rep gene, complete cap gene and 3' ITR is 98% identical to AAV-1.

Biologically, such recombination may enable AAV-1 to acquire the ability to transmit through the human population. It is also interesting to note that the ITRs of AAV-6 comprise one AAV-1 ITR and one AAV-2 ITR. The replication model of defective parvovirus can maintain this special arrangement. Studies on AAV integration have shown that a majority of AAV integrants carries deletions in at least one of the terminal repeats. These deletions have been shown to be able to be repaired through gene conversion using the other intact terminal repeat as a template. Therefore, it would be very difficult to maintain AAV-6 as a homogenous population

when an integrated copy of AAV-6 is rescued from host cells with helper virus infection. The AAV-6 with two identical AAV-2 ITRs or two identical AAV-1 ITRs should be the dominant variants. The AAV-6 with two AAV-1 ITRs has been observed by Russell's group [Rutledge, cited above (1998)]. So far there is no report on AAV-6 with two AAV-2 ITRs. Acquirement of AAV-2 P5 promoter by AAV-6 may have explained that AAV-6 have been isolated from human origin while AAV-1 with the same virion has not. The regulation of P5 promoter between different species of AAV may be different *in vivo*. This observation suggests the capsid proteins of AAV were not the only determinants for tissue specificity.

10

15

5

Although it is clear that AAV-6 is a hybrid of AAV-1 and AAV-2, AAV-6 has already exhibited divergence from either AAV-1 or AAV-2. There are two nucleotide differences between AAV-6 and AAV-2 in their first 450 nucleotides. There are about 1% differences between AAV-6 and AAV-1 in nucleotide levels from nucleotides 522 to the 3' end. There also exists a quite divergent region (nucleotide 4486-4593) between AAV-6 and AAV-1 (Figs. 1A-1C). This region does not encode any known proteins for AAVs. These differences in nucleotide sequences may suggest that AAV-6 and AAV-1 have gone through some evolution since the recombination took place. Another possible explanation is that there exists another variant of AAV-1 which has yet to be identified. So far, there is no evidence to rule out either possibility. It is still unknown if other hybrids (AAV-2 to AAV-4, etc.) existed in nature.

20

The coding region of AAV-1 was deduced by comparison with other known AAV serotypes. Table 1 illustrates the coding region differences between AAV-1 and AAV-6. The amino acid residues are deduced according to AAV-2.

25

With reference to the amino acid position of AAV-1, Table 1 lists the amino acids of AAV-1 which have been changed to the corresponding ones of AAV-6. The amino acids of AAV-1 are shown to the left of the arrow. Reference may be made to SEQ ID NO: 5 of the amino acid sequence of AAV-1 Rep 78 and to SEQ ID NO: 13 for the amino acid sequence of AAV-1 VP1.

20

24

Table 1
Coding region variations between AAV-1 and AAV-6

Rep protein (Rep78)			Cap protein (VP1)	
Position(s)	Amino acids		Position(s)	Amino acids
28	S-N.	]	129	L→F
191	Q-H	.	418	· E-D
192	. H-D		531	E-K
308	E-D		584	F→L
			598	A→V
			642	N→H

It was surprising to see that the sequence of the AAV-1 coding region is almost identical to that of AAV-6 from position 452 to the end of coding region (99%). The first 508 nucleotides of AAV-6 have been shown to be identical to those of AAV-2 [Rutledge, cited above (1998)]. Since the components of AAV-6 genome seemed to be AAV-2 left ITR - AAV-2 p5 promoter - AAV-1 coding region - AAV-1 right ITR, it was concluded that AAV-6 is a naturally occurred hybrid between AAV-1 and AAV-2.

## Example 4 - Gene Therapy Vector Based on AAV-1

Recombinant gene transfer vectors based on AAV-1 viruses were constructed by the methods described in Example 1. To produce a hybrid recombinant virus with AAV-1 virion and AAV-2 ITR, the AAV-1 trans plasmid (pAV1H) and the AAV-2 cis-lacZ plasmid (with AAV-2 ITR) were used. The AAV-2 ITR was used in this vector in view of its known ability to direct site-specific integration. Also constructed for use in this experiment was an AAV-1 vector carrying the green fluorescent protein (GFP) marker gene under the control of the immediate early promoter of CMV using pAV1H as the trans plasmid.

20

25

25

## A. rAAV-1 Viruses Transfect Host Cells in Vitro

84-31 cells, which are subclones of 293 cells (which express adenovirus E1a, E1b) which stably express E4/ORF5, were infected with rAAV-1 GFP or rAAV-lacZ. High levels of expression of GFP and lacZ was detected in the cultured 84-31 cells. This suggested that rAAV-1 based vector was very similar to AAV-2 based vectors in ability to infect and expression levels.

#### B. <u>rAAV-1 Viruses Transfect Cells in Vivo</u>

The performance of AAV-1 based vectors was also tested *in vivo*. The rAAV-1 CMV-α1AT virus was constructed as follows. The EcoRI fragment of pAT85 (ATCC) containing human α1-antitrypsin (α1AT) cDNA fragment was blunted and cloned into PCR (Promega) at a Smal site to obtain PCR-α1AT. The CMV promoter was cloned into PCR-α1AT at the Xbal site. The Alb-α1AT expression cassette was removed by XhoI and ClaI and cloned into pAV1H at the Xbal site. This vector plasmid was used to generate AAV-1-CMV-α1AT virus used in the experiment described below.

For screening human antibodies against AAV, purified AAV virus is lysed with Ripa buffer (10 mM Tris pH 8.2, 1% Triton X-100, 1% SDS, 0.15 M NaCl) and separated in 10% SDS-PAGE gel. The heat inactivated human serum was used at a 1 to 1000 dilution in this assay. The rAAV-1 CMV-α1AT viruses were injected into Rag-1 mice through tail vein injection at different dosages. The concentration of human α1-antitrypsin in mouse serum was measured using ELISA. The coating antibody is rabbit anti-human human α1-antitrypsin (Sigma). The goatantihuman α1-antitrypsin (Sigma) was used as the primary detection antibodies. The sensitivity of this assay is around 0.3 ng/ml to 30 ng/ml. The expression of human α-antitrypsin in mouse blood can be detected in a very encouraging level. This result is shown in Table 2.

PCT/US99/25694

5

15

20

25

26

Table 2
Human Antitrypsin Expressed in Mouse Liver

Amount of virus injected	Week 2 (ng/ml)	Week 4 (ng/ml)
2x10 <sup>10</sup> genomes	214.2	171.4
1x10 <sup>10</sup> genomes	117.8	109.8
5x10 <sup>10</sup> genomes	64.5	67.8
2.5x10 <sup>10</sup> genomes	30.9	58.4

rAAV-1 CMV-lacZ viruses were also injected into the muscle of
C57BL6 mice and similar results were obtained. Collectively, these results suggested
that AAV-1 based vector would be appropriate for both liver and muscle gene
delivery.

#### Example 5 - Neutralizing Antibodies Against AAV-1

Simple and quantitative assays for neutralizing antibodies (NAB) to AAV-1 and AAV-2 were developed with recombinant vectors. A total of 33 rhesus monkeys and 77 normal human subjects were screened.

#### A. Nonhuman Primates

Wild-caught juvenile rhesus monkeys were purchased from Covance (Alice, Tex.) and LABS of Virginia (Yemassee, SC) and kept in full quarantine. The monkeys weighed approximately 3 to 4 kg. The nonhuman primates used in the Institute for Human Gene Therapy research program are purposefully bred in the United States from specific-pathogen-free closed colonies. All vendors are US Department of Agriculture class A dealers. The rhesus macaques are therefore not infected with important simian pathogens, including the tuberculosis agent, major simian lentiviruses (simian immunodeficiency virus and simian retroviruses), and cercopithecine herpesvirus. The animals are also free of internal and external parasites. The excellent health status of these premium animals minimized the potential for extraneous variables. For this study, serum was obtained from monkeys prior to initiation of any protocol.

10

15

20

25

NAB titers were analyzed by assessing the ability of serum antibody to inhibit the transduction of reporter virus expressing green fluorescent protein (GFP) (AAV1-GFP or AAV2-GFP) into 84-31 cells. Various dilutions of antibodies preincubated with reporter virus for 1 hour at 37°C were added to 90% confluent cell cultures. Cells were incubated for 48 hours and the expression of green fluorescent protein was measured by FluoroImaging (Molecular Dynamics). NAB titers were calculated as the highest dilution at which 50% of the cells stained green.

Analysis of NAB in rhesus monkeys showed that 61% of animals tested positive for AAV-1; a minority (24%) has NAB to AAV-2. Over one-third of animals had antibodies to AAV-1 but not AAV-2 (i.e., were monospecific for AAV-1), whereas no animals were positive for AAV-2 without reacting to AAV-1. These data support the hypothesis that AAV-1 is endemic in rhesus monkeys. The presence of true AAV-2 infections in this group of nonhuman primates is less clear, since cross-neutralizing activity of an AAV-1 response to AAV-2 can not be ruled out. It is interesting that there is a linear relationship between AAV-2 NAB and AAV-1 NAB in animals that had both.

#### B. Humans

For these neutralization antibody assays, human serum samples were incubated at 56°C for 30 min to inactivate complement and then diluted in DMEM. The virus (rAAV or rAd with either lacZ or GFP) was then mixed with each serum dilution (20X, 400X, 2000X, 4000X, etc.) and incubated for 1 hour at 37°C before applied to 90% confluent cultures of 84-31 cells (for AAV) or Hela cells (for adenovirus) in 96-well plates. After 60 minutes of incubation at culture condition, 100 µl additional media containing 20% FCS was added to make final culture media containing 10% FCS.

The result is summarized in Table 3.

Table 3

Adenovirus	AAV-1	AAV-2	# of samples	Percentage
•	-	-	41	53.2%
+	-		16	20.8%
•	+	-	0	0.0%
-	-	+	2	2.6%
-	+	+	2	2.6%
+	_	.+	3	3.9%
+	+	-	0	0.0%
+	+	+	13	16.9%
		Total	77	100%

10

15

20

25

5

The human neutralizing antibodies against these three viruses seemed to be unrelated since the existence of neutralizing antibodies against AAV are not indications for antibodies against adenovirus. However, AAV requires adenovirus as helper virus, in most of the cases, the neutralizing antibodies against AAV correlated with the existence of neutralizing antibodies to adenovirus. Among the 77 human serum samples screened, 41% of the samples can neutralize the infectivity of recombinant adenovirus based on Ad5. 15/77 (19%) of serum samples can neutralize the transduction of rAAV-1 while 20/77 (20%) of the samples inhibit rAAV-2 transduction at 1 to 80 dilutions or higher. All serum samples positive in neutralizing antibodies for AAV-1 in are also positive for AAV-2. However, there are five (6%) rAAV-2 positive samples that failed to neutralize rAAV-1. In samples that are positive for neutralizing antibodies, the titer of antibodies also varied in the positive ones. The results from screening human sera for antibodies against AAVs supported the conclusion that AAV-1 presents the same epitome as that of AAV-2 to interact

10

15

20

25

with cellular receptors since AAV-1 neutralizing human serums can also decrease the infectivity of AAV-2. However, the profile of neutralizing antibodies for these AAVs is not identical, there are additional specific receptors for each AAV serotype.

## Example 6 - Recombinant AAV Viruses Exhibit Tissue Tropism

The recombinant AAV-1 vectors of the invention and the recombinant AAV-2 vectors [containing the gene encoding human α1-antitrypsin (α1AT) or murine erythropoietin (Epo) from a cytomegalovirus-enhanced β-actin promoter (CB)] were evaluated in a direct comparison to equivalent copies of AAV-2 vectors containing the same vector genes.

Recombinant viruses with AAV-1 capsids were constructed using the techniques in Example 1. To make rAAV with AAV-1 virions, pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide Rep and Cap functions. For the generation of the rAAV based on AAV-2, p5E18(2/2) was used as the *trans* plasmid, since it greatly improved the rAAV yield. [Early experiments indicated similar *in vivo* performances of AAV-1 vectors produced with pAV1H and p5E19 (2/1). All subsequent studies used AAV-1 vectors derived from p5E18(2/1) because of the increased yield.]

Equivalent stocks of the AAV-1 and AAV-2 vectors were injected intramuscularly (5 x 10<sup>10</sup> genomes) or liver via the portal circulation (1 x 10<sup>11</sup> genomes) into immunodeficient mice, and the animals (four groups) were analyzed on day 30 for expression of transgene. See, Figs. 4A and 4B.

AAV-2 vectors consistently produced 10- to 50-fold more serum erythropoietin or a1-antitrypsin when injected into liver compared to muscle. (However, the AAV-1-delivered genes did achieve acceptable expression levels in the liver.) This result was very different from that for AAV-1 vectors, with which muscle expression was equivalent to or greater than liver expression. In fact, AAV-1 outperformed AAV-2 in muscle when equivalent titers based on genomes were administered.

#### Example 7 - Gene Delivery via rAAV-1

5

10

15

20

25

30

C57BL/6 mice (6- to 8-week old males, Jackson Laboratories) were analyzed for AAV mediated gene transfer to liver following intrasplenic injection of vector (i.e., targeted to liver). A total of 10<sup>11</sup> genome equivalents of rAAV-1 or rAAV-2 vector were injected into the circulation in 100 μl buffered saline. The first vector contained either an AAV-1 capsid or an AAV-2 capsid and expressed α1AT under the control of the chicken β-actin (CB) promoter. Day 28 sera were analyzed for antibodies against AAV-1 or AAV-2 and serum α1AT levels were checked. Animals were then injected with an AAV-1 or AAV-2 construct expressing erythropoietin (Epo, also under the control of the CB promoter). One month later sera was analyzed for serum levels of Epo. The following groups were analyzed (Figs. 5A-5D).

In Group 1, vector 1 was AAV-2 expressing a1AT and vector 2 was AAV-2 expressing Epo. Animals generated antibodies against AAV-2 following the first vector administration which prevented the readministration of the AAV-2 based vector. There was no evidence for cross-neutralizing the antibody to AAV-1.

In Group 2, vector 1 was AAV-1 expressing a1AT while vector 2 was AAV-1 expressing Epo. The first vector administration did result in significant a1AT expression at one month associated with antibodies to neutralizing antibodies to AAV-1. The animals were not successfully readministered with the AAV-1 Epo expressing construct.

In Group 3, the effectiveness of an AAV-2 vector expressing Epo injected into a naive animal was measured. The animals were injected with PBS and injected with AAV-2 Epo vector at day 28 and analyzed for Epo expression one month later. The neutralizing antibodies were evaluated at day 28 so we did not expect to see anything since they received PBS with the first vector injection. This shows that in naive animals AAV-2 is very efficient at transferring the Epo gene as demonstrated by high level of serum Epo one month later.

Group 4 was an experiment similar to Group 3 in which the animals originally received PBS for vector 1 and then the AAV-1 expressing Epo construct 28 days later. At the time of vector injection, there obviously were no antibodies to either

10

15

20

25

AAV-1 or AAV-2. The AAV-1 based vector was capable of generating significant expression of Epo when measured one month later.

Group 5 is a cross-over experiment where the initial vector is AAV-2 expressing α1AT followed by the AAV-1 construct expressing Epo. The animals, as expected, were efficiently infected with the AAV-2 vector expressing α1AT as shown by high levels of the protein in blood at 28 days. This was associated with significant neutralizing antibodies to AAV-2. Importantly, the animals were successfully administered AAV-1 following the AAV-2 vector as shown by the presence of Epo in serum 28 days following the second vector administration. At the time of this vector administration, there was high level AAV-2 neutralizing antibodies and very low cross-reaction to AAV-1. The level of Epo was slightly diminished possibly due to a small amount of cross-reactivity. Group 6 was the opposite cross-over experiment in which the initial vector was AAV-1 based, whereas the second experiment was AAV-2 based. The AAV-1 vector did lead to significant gene expression of α1AT, which also resulted in high level AAV-1 neutralizing antibody. The animals were very efficiently administered AAV-2 following the initial AAV-1 vector as evidenced by high level Epo.

A substantially identical experiment was performed in muscle in which 5 x 10<sup>10</sup> genomes were injected into the tibialis anterior of C57BL/6 mice as a model for muscle directed gene therapy. The results are illustrated in Figs. 6A-6D and are essentially the same as for liver.

In summary, this experiment demonstrates the utility of using an AAV-1 vector in patients who have pre-existing antibodies to AAV-2 or who had initially received an AAV-2 vector and need readministration.

#### Example 8 - Construction of Recombinant Viruses Containing AAV-1 ITRs

This example illustrates the construction of recombinant AAV vectors which contain AAV-1 ITRs of the invention.

An AAV-1 cis plasmid is constructed as follows. A 160 bp Xho-NruI AAV-1 fragment containing the AAV-1 5' ITR is obtained from pAV1-BL. pAV1-BL was

10

15

20

25

generated as described in Example 1. The Xho-NruI fragment is then cloned into a second pAV1-BL plasmid at an XbaI site to provide the plasmid with two AAV-1 ITRs. The desired transgene is then cloned into the modified pAV-1BL at the NruI and BamHI site, which is located between the AAV-1 ITR sequences. The resulting AAV-1 cis plasmid contains AAV-1 ITRs flanking the transgene and lacks functional AAV-1 rep and cap.

Recombinant AAV is produced by simultaneously transfecting three plasmids into 293 cells. These include the AAV-1 cis plasmid described above; a trans plasmid which provides AAV rep/cap functions and lacks AAV ITRs; and a plasmid providing adenovirus helper functions. The rep and/or cap functions may be provided in trans by AAV-1 or another AAV serotype, depending on the immunity profile of the intended recipient. Alternatively, the rep or cap functions may be provided in cis by AAV-1 or another serotype, again depending on the patient's immunity profile.

In a typical cotransfection, 50 µg of DNA (cis:trans:helper at ratios of 1:1:2, respectively) is transfected onto a 15 cm tissue culture dish. Cells are harvested 96 hours post transfection, sonicated and treated with 0.5% sodium deoxycholate (37° for 10 min). Cell lysates are then subjected to 2-3 rounds of ultracentrifugation in a cesium gradient. Peak fractions containing rAAV are collected, pooled and dialyzed against PBS. A typical yield is 1 x 10<sup>13</sup> genomes/10<sup>9</sup> cells.

Using this method, one recombinant virus construct is prepared which contains the AAV-1 ITRs flanking the transgene, with an AAV-1 capsid. Another recombinant virus construct is prepared with contains the AAV-1 ITRs flanking the transgene, with an AAV-2 capsid.

All publications cited in this specification are incorporated herein by reference. While the invention has been described with reference to a particularly preferred embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the claims.

#### What is claimed is:

- 1. An isolated AAV-1 nucleic acid molecule comprising a sequence selected from the group consisting of:
  - (a) SEQ ID NO: 1;
  - (b) a DNA sequence complementary to SEQ ID NO: 1;
  - (c) cDNA complementary to (a) or (b); and
  - (d) RNA complementary to any of (a) to (c).
- 2. A nucleic acid molecule comprising an AAV-1 inverted terminal repeat (ITR) sequence selected from the group consisting of:
  - (a) nt 1 to 143 of SEQ ID NO: 1;
  - (b) nt 4576 to 4718 of SEQ ID NO: 1;
  - (c) a nucleic acid sequence complementary to (a) or (b); and
  - (d) a functional fragment of (a), (b), or (c).
- 3. A recombinant vector comprising a 5' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
  - (a) nt 1 to 143 of SEQ ID NO: 1;
  - (b) a nucleic acid sequence complementary to (a); and
  - (c) a functional fragment of (a) or (b).
- 4. The recombinant vector according to claim 3, wherein said vector further comprises a 3' AAV-1 ITR.

- 5. A recombinant vector comprising a 3' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
  - (a) nt 4576 to 4718 of SEQ ID NO: 1;
  - (b) a nucleic acid sequence complementary to (a); and
  - (c) a functional fragment of (a) or (b).
- 6. The recombinant vector according to claim 5, wherein said vector further comprises a 5' AAV-1 ITR.
- 7. The recombinant vector according to any of claims 3-6, wherein said vector further comprises AAV-1 capsid proteins having the sequence of SEQ ID NO: 13, 15 or 17 or functional fragments thereof.
- 8. The recombinant vector according to any of claims 3-6, wherein said vector further comprises adenovirus sequences.
- 9. A recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.
- 10. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said cap coding region comprises at least one member is selected from the group consisting of:
  - (a) vp1, nt 2223 to 4431 of SEQ ID NO: 1;
  - (b) vp2, nt 2634 to 4432 of SEQ ID NO: 1; and
  - (c) vp3, nt 2829 to 4432 of SEQ ID NO: 1.

- 11. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said rep coding region comprises an AAV-1 rep coding region comprising at least one member selected from the group consisting of:
  - (a) rep 78, nt 335 to 2304 of SEQ ID NO: 1;
- (b) rep 68, nt 335 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto,
  - (c) rep 52, nt 1007 to 2304 of SEQ ID NO: 1; and
- (d) rep 40, nt 1007 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto.
- 12. A host cell transduced with a recombinant viral vector according to any of claims 3-6.
- 13. A host cell transduced with a nucleic acid molecule according to any of claims 1, 2, 10 or 11.
- 14. A host cell stably transduced with an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1.
- 15. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to any of claims 3-6.
- 16. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 7.
- 17. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 8.

- 18. A method for AAV-mediated delivery of a transgene comprising the step of delivering to a host cell an AAV virion which comprises:
- (a) a capsid comprising at least one capsid protein encoded by an AAV-1 cap gene; and
- (b) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 19. A method for AAV-mediated delivery of a transgene to a host comprising the steps of:
- (a) assaying a sample from the host to determine the presence of neutralizing antibodies specific against any serotype of AAV; and
  - (b) delivering to the host an AAV virion which comprises:
- (i) a capsid comprising at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has no antibodies as determined in step (a); and
- (ii) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 20. The method according to claim 19, comprising the additional step of repeating steps (a) and (b).
- 21. Use of an AAV virion which comprises a capsid comprising (a) at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has antibodies, and (b) a DNA molecule comprising a transgene operably linked to regulatory sequences directing its expression,

in the preparation of a medicament for delivery of a transgene to a host, wherein said host has no preexisting neutralizing antibodies against the AAV serotype of said cap gene.

37

- 22. A method for delivery of a transgene comprising the step of delivering to a host cell a recombinant virus comprising a recombinant vector according to any of claims 3-8.
- 23. A method for producing a selected gene product comprising the steps of transfecting a mammalian cell with the molecule according to claim 1 or a functional fragment thereof and culturing said cell under conditions suitable to express said gene product.

## FIG IA

	Neb amanig area	120
AAV-1	ttgeccaetecetetetgegegetegetegeteggtggggcetgeggaecaaaggteegeagaeggeagagetetgetet	120
AAV-2	tigeceaticcitatigegogategategategategategategategategategate	120
AAV-8	a dien	
	TRS <u>E box/USF</u> **greeneteenetergyyytentcoccaaccccccccccccccccccccccccaccctaacctaactaactaactacaccac	
AAV-1	ggesafteentesetaggggtantegegaagegeeteeegegegegegegegegaegtaagtaa	337
AAV-3	ggcaactccascatggygrandcacatacatacatacatacatacatacatacatacatac	222
AAV-6	.c	
	YYI PS/TATA YYI/oS RNA Rep78/68	
	THE PARTY OF THE P	356
AAV-2	GACTITICCACT T. A.C. A.C. G. AG. G. G. G. G. G. G. T. T. A.C. T. A.C. AC. G. AG. G. G. G. G. G. G. G. T. AT. G. T. A.C. AC. G. AG. G. G. G. G. G. G. G. T. AC. T. AC. AC. G. AG. G. G. G. G. G. G. G. G. T. AC. T. AC. AC. G. AG. G. G. G. G. G. G. G. G. G. T. AC. T. AC. T. AC. AC. G. AG. G. G. G. G. G. G. G. G. G. T. AC. T. AC. T. AC. G. AG. G. T. AC. T. AC. G. AC. G. AG. G. G	343
AAV-6		341
	TEATCAAGCTGCCGAGCGACCTGCAGGGCACCTGCCGGGCATTTCTGACTCGTTTGTGACCTGGCTGG	476
		***
AAV - 3	TCTTTAGCA	462
~~		
	· ·	•
AAV-1	TTELECACCECCETCACCOTCOCCEACAACCTCCACCCCCACTTCCTGOTCCAATCCCCCCCTCACTAACCACCCCCCCTCATCTTTCTTC	596
	T 1000 m 'T	304
AAV-6		347
	ACTICEACTICATA TICTOUTOGNOACCACOGGOGTCANATCCATOGTOCTOCGCCCCTTCLTCAGTCAGTCAGATAGGGACAACCTGGTGCAGACCATCTACCGCGGGATGGAGCCGACCC	716
	1 0 00 0 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1	,
AAV-6		701
LAV-1	TECCCALCITECTICE COGTOLCCALGACOCOTAL TEGEOGCICGALOCCECCALCIACOTOSTOCALCALGIOSTOCIACA TECCCALCIACIAC TECCCALGACIAC COCCALGACIAC COCALGACIAC COCCALGACIAC COCCALGACIAC COCCALGACIAC COCCALGACIAC COCA	122
MAA-3		621
	P19/TATA P19 RNA COTOGRATALATOGRAGIATATALOGOCICTOTTTCALCITGOCCCALOGECALOCACCACCACCACCACCACCACCACCACCACCACCACCAC	
	CONTROL OF THE PRODUCT OF THE PRODUC	956
MV-6		941
	Rep 52/40 ACCCCAATTCTGACGCCCCTGTCATCCGCTCAAAAACCTCCGGCCCCTACATGGGCTGGCT	1076
MV-1	.T	1062
TVA-3		1061
WA-8		
AV-1	CCTCGTACATCTCCAACCCCCCTTCCAACTCCCACTCCCAGATCAAGGCCCCTTCGGACAATGCCGGCAAGATCATGCCGCCAACTACCTGGTACCAACTCAGCACTACCTGGTACGACTAACCTGGTACACACTAACCTGGTACACACAC	1170
WA-3		1101
MA-2		
•		
NW-1	CTCCGCCCCCCCCCACATTALAACCAACCCCATCTACCGCATCCTGCACCTGCACCTTACCACCTACCCCTACCCTCCCT	1316
	- LO - COMPO L	
AV-6		1301
		****
1-Va	ACACCAMETROCTOTTTCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	1422
YA-3		1421
74-6		•
AV. 1	ATTOCOTCOACAAGATOOTGATCTGGTGGGGGGGGGGGCAAGATGACGCCCAAGGTCGTGGAGTCCGCCAAGGCCATTCTCGGGCGCCAGCAAGGTGCCCAAAAGTGCAAGTCGT	2556
		T367
14-6		1341
7A-1	CCCCCCACATCGACCCCCCCCCCCCCCCCCCCCCCCCCC	1667
7A - 3	.GAG.T	1661
AV-6	<del></del>	<del>-</del>
AV.1	TTGLACTCACCCCCCTTTGGLACCATGACTTTGGCLAGGTGACAAAGCAGGAAGTCAAGTC	1796
1	Ψ	1104
AV - 4		1781
	P40/TATA P40 RNA	
W-1 1	ACCOTOGRACIA MANCA COCCCCCCC CONTENA COCCCCATAMA O CONSCICA MACCOCCCC COCCCCC COCCCC COCCC COCCCC COCCCCCC	1110
L- V	AGGSTGRAFTARCALTCACCCCCCCCATCACCCCATCACCCCACCCCCCCCC	2477
	##	

## FIG 1B

	1 CCCLCLOGTACCAAACAATGTTCTCGTCACGCGGGCATGCTTCAGATGCTGTTTCCCTGCLAGACATGCGAGAAATGAATGAATGAATTTCACATTTGCTTCACGCACG	
AAV-2	2 <u>,AChChCh</u>	. 3019
WA-	f ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	. 2021
	·	
	A CTUTTCHEAGTGCTTCCCCCCCCTTCCAGAATCTCAACCGTTCGTCAGAAAAACGACGTATCGGAAACTCTGTGCCATTCATCATCATCTCCTCGGGGGGGCTCCCCGAGATTCATTC	P 2151
AAV.	T	2111
AAV.	A	. 2131
	Rep78 stop VP1 Technologic Rep68 stop VP1 Rep68 stop VP1 Technologic Rep68	stop
AAV-1	COGCCTGCGATCTGGTCAACGTGGACGGGGGGGGGGCTGTTTCTGAGCAATAAATGACTTAAACGAGGTATGGCTGCCGATGGTTATCTTCCAGAATGGCTGGAGGACAACCTCTCTGAG	2271
AAV-3		2253
77A-6	; ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	2254
110.1	oacttochescetcharitaliactecharitaliacepocologicochiscalcalicalia and careeal a	2191
MY-3	.A., AA, AC, A. GC. C CC. A ACCA A GC GCAG GGC. TA	2373
AAV-6		2378
	•	
WAA-J	GENETICALENAGGGGAGCCCGTCNAGGCGGGGGGGGGGGGGGGGGGGGG	2513
AAV-2	AGAAG	2493
YYA-2	h.000T	2478
AAV-1	GCCCAOTTTCAOGAGCOTCTOCAAGAAGATACOTCTTTTGGGGGCAACCTCGGGCGAAGATCTCCAGCCAAGAAGCGGGTTCTCGAACCTCTCGCTCTGAGGAAGGCGCTAAG	2633
AAV-2		2613
AAV- F	C	2610
	A188	
	<u>VP2</u>	
AAV-1	ACCUPATION AND AND AND AND AND AND AND AND AND AN	2733
AAV-3	GAGA.G	2733
~~~		
	VP3	
444-1	GACTCAGAGTCAGTCCCCGATCCACAACCTCTCGGAGAACCTCCAGCAACCCCCGCTGCTGTGACCACTACTACAATGGCTTCAGGCGCTGGCGCACCAATGGCAGACAATAACGAAGGC	2873
AA7-2	GCATCCGGG.GAG.GTG.CAAGAAAAA	2853
<b>447-6</b>	<del></del>	2158
	GCCGACGCGATGCGTAATCCCTCAGGAAATTGCCATTCCGATTCCCACTGCCCGACGGCGACGGACAGAGTCATCACCACCCAC	2002
114-2	T. C.	2971
AAV-6		2978
	AMCANATOTOCHGTECTTCHACGGGGGGCCAGCACGACAACCACTACTTCGGCTACAGCACCCCCGGGGGGTATTTCATTTCAACAGATTCCACTGGCACTTTTCACCAGGACTGG	
	ATCCAAATCGTTT	
VTA-2		3090
117.1	CAGGGACTEA TEANCALCART TECOGGATTECOGGCCCAAGAGACTCAACTTCAACTTCAACTTCAACTAGGACGCTTCACGACCAATCATCATCATCATCATCATCATCATCATCATCATC	1211
	.AA	
147-6	g	3218
	••	
WA-7	ACCAMINATION AND TOTAL TOTAL TOTAL TRANSPORTATION AND AND AND AND AND AND AND AND AND AN	3353
WA-3		3330
VAA-6		3338
127-1	TACCTGACGCTCAACAATGGCGACCCAAGCCGTGGGACCTTCATCCTTTTACTGCCTGGGATATTTCCCTTCTCAGAACGCGCGACAACAACTTTACCTTCAGCTACACCTTTCAG	3473
	C. G. G C G T G. A. A. A G. T. A	
74-6		3438
	•	
WA-1	GALGTOCCTTTCCACAGCAGCTACGGGCACAGCCAGCCCTGACGGCTGATGAATCCTCTCATCGACCAATACCTGTATTACCTGAACAGAACTCAAAATCAGTCCCAAATCCCCAA	3593
74.3		3570
MA-R	C	3378
AV-1	ANCANGIAC TROUTE TARGET TO THE CONTROL TO THE CONTR	3711
WA-1	C.GTCARGCC.T.A. TCT.AG.COGAG.GAG.A. TCGG.AC. T.T.GG	3690
MV-4	G	3694
	. ,	
AV-1	ANCACCANTITIACCTCCACCTCCTTCAAAATATAACCTCAATCCCCTCATCA	3833
74-3	TG.A.AET.GAA.CGGCCCCA.ACTC.GG.GTGGC.CCAAGCGTATT	3010

## FIG IC

AAV-2	AGCGGTGTCATGATTTTTGGAAAAGAGAGGGGCGCGGAGCTTCAAACACTGCATTGGACAATGTCATGATTACAGAGGAAGAGGAAATTAAAGCCACTAACCCTGTGGCCACCGAAAAATTACAGACGAAGAAGAATTAAAGCAGAAAAACC.TT.G.,GCAGAA.G.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAGAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAGAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,GCAG,A.,G	3930
	OGGACCGTGGCAGTCAATTTCCAGAGCAGCACAGCACCCTGCAACCCTGCAGATGTGCATGCTATGCAAGCATTACCTCGCATGGTUTGGCAAGAATAGAGAAGTGTAACCTGCAGGGTTACCT	
AAV-1	ATTTGGGCCAAAATTCCTCACACAGATGGACACTTTCACCCGTCTCTCTTATGGGCGGCTTTGGACTCAAGAACCGGCCTCCTCAGATCCTCAGATCCTCAAAAAACACGGCCTUTTCCTGGGAAT	4193
	.chGAGCTCCCCThThTGCGhT	
	CCTCCGGCGGAGTTTCAGCTACAAAGTTTGCTTCATTCAT	
	TA.CACCCAGTGG	
LAV-1	CCAACTGCAGTACACATCAATTATGCAAAATCTGCCCAA-CGTTCATTTTACTGTACAACAACAATGCACTTTATATATTATCCACCCCCATTACCATTACCTTACCCTTACCCTTACCACC	4431
WA-3		.1410
	VP1-3 stop PolyA signal	
LAV-1	ANTIAL GROUP TRANSCRIPT ANTICATION TO THE CAST OF THE	4547
WA-3	G.TT	4530
WA- 6	G7	4533
AV-1	TTCCCCATALAACACTTACGTCATCCCCCttacccctagtgatggagttgcccactccctctctgcgcgctcggtcgg	4667
AV-2	ACTA.A.gg.agc.c.a.gc.gc.c.a.gc.c.gca.gc.c.gg	4630
47-6	At.	4632
	Egytocycagycoccaccyagcyagcycycocagagaggyagtgggcaa 471%	
	ac.g.gatgt	
7A-Q	1683	

--aav-2 itr --AAV-1 ITR -aav-1 T A
C G
GC Cg
C G
A Tg a Ca
GCGGGGTGGCTCGCTCCCTCACCCGTT FIG 2

AAV-1 TR




FIG. 3A




FIG. 3B







FIG. 4A





Group	1	2	3	4	5	6
Vector1 - a1AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2



Group	1	2	3	4	5	6
Vector1 - a1AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2

## SEQUENCE LISTING

```
<110> Wilson, James M.

Xiao, Weidong

The Trustees of the University of Pennsylvania
```

<120> Adeno-Associated Virus Serotype I Nucleic Acid Sequences, Vectors and Host Cells Containing Same

<130> GNVPN.031PCT

<140>

<141>

<150> 60/107,114

<151> 1998-11-05

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 4718

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (335)..(2206)

· <220>

<221> CDS

<222> (2223)..(4430)

<400> 1

ttgcccactc cctctctgcg cgctcgctcg ctcggtgggg cctgcggacc aaaggtccgc 60

agacggcaga gctctgctct gccggcccca ccgagcgagc gagcgcgcag agagggagtg 120

ggcaactcca tcactagggg taatcgcgaa gcgcctccca cgctgccgcg tcagcgctga 180

cgtaaattac gtcatagggg agtggtcctg tattagctgt cacgtgagtg cttttgcgac 240

attttgcgac accacgtggc catttagggt atatatggcc gagtgagcga gcaggatctc 300

cattttgacc gcgaaatttg aacgagcagc agcc atg ccg ggc ttc tac gag atc 355

Met Pro Gly Phe Tyr Glu Ile

gtg	atc	aag	gtg	ccg	agc	gac	ctg	gac	gag	cac	ctg	ccg	ggc	att	tct	403
Val	Ile	Lys	Val	Pro	Ser	Asp	Leu	Asp	Glu	His	Leu	Pro	Gly	Ile	Ser	
		10					15					20				
gac	tcg	ttt	gtg	agc	tgg	gtg	gcc	gag	aag	gaa	tgg	gag	ctg	ccc	ccg	451
Asp	Ser	Phe	Val	Ser	Trp	Val	Ala	Glu	Lys	Glu	Trp	Glu	Leu	Pro	Pro	
	25					30					35					
gat	tct	gac	atg	gat	ctg	aat	ctg	att	gag	cag	gca	ccc	ctg	асс	gtg	499
Asp	Ser	Asp	Met	Asp	Leu	Asn	Leu	Ile	Glu	Gln	Ala	Pro	Leu	Thr	Val	
40					45					50					55	
gcc	gag	aag	ctg	cag	cgc	gac	ttc	ctg	gtc	caa	tgg	cgc	cgc	gtg	agt	547
Ala	Glu	Lys	Leu	Gln	Arg	Asp	Phe	Leu	Val	Gln	Trp	Arg	Arg	Val	Ser	
				60					65					70		
														•		
aag	gcc	ccg	gag	gcc	ctc	ttc	ttt	gtt	cag	ttc	gag	aag	ggc	gag	tcc	595
Lys	Ala	Pro	Glu	Ala	Leu	Phe	Phe	Val	Gln	Phe	Glu	Lys	Gly	Glu	Ser	
_			75					80				_	85			
tac	ttc	cac	ctc	cat	att	ctq	qtq	gaq	acc	acq	aaa	qtc	aaa	tcc	atq	643
															Met	
-,-		90					95				2	100	_,			
ata	cta	aac	cac	ttc	ctg	agt	caq	att	agg	gac	aaq	cta	ata	caσ	acc	691
	_		_		Leu						_	_		_		
	105	•	_			110			,	•	115	•				
atc	tac	cqc	ggg	atc	gag	ccg	acc	ctq	ccc	aac	tqq	ttc	qcq	ata	acc	739
		-			Glu	_		_								
120	-		-		125					130	•				135	
aaq	acq	cat	aat	aac	acc	qqa	aaa	aga	aac	aaq.	ata	ata	gac	gag	tgc	787
-	_	-			Ala					-				-	-	
• •		•		140		•	•	-4	145	•				150	-,-	
tac	atc	ccc	aac	tac	ctc	cta	ccc	aaσ	act	cad	ccc	gag	cta	cag	taa	835
					Leu	_		-		_				_		
- , -			155	- 1 -				160					165	·		
				٠											•	
aca	taa	act	aac	ato	gag	gag	tat	ata	age	acċ	tat	tta	aac	cta	acc	883
											_	-		_	Ala	0,00
n.a	1 - P	170	7311	.10 L	JIU	Jiu	175	116	JUL	7.1 a	Uys	180	usii	TEU	VT a	
		170					1/3					100				
~~~	cac	222	caa	ctc	gtg	aca	C= 4	cac	cta			ata		<b>~=</b> ~	300	021
					Val	-		•				_	_			931
GIU	185	пλэ	ALG	Ten	val	190	GTII	117.2	ъец	TIIL		val	Set	GIII	1111	
	707					170					195					

		cag Gln														979
		tca Ser					-		-		-	-			-	1027
, ,	-	cgg Arg						-					, ,	•	_	1075
_	_	tac Tyr 250					-	_			-			_		1123
-	-	gct Ala	_	-		-		-		-		_				1171
		gac Asp		_	-			-	-			-				1219
	-	atc Ile		-		_		-				-		-		1267
-	-	tcc Ser	_					•	_					_	_	1315
		atc Ile 330		-			_	-		•		-				1363
	-	gcc Ala		_		-						-	-			1411
		gag Glu						_	-	•	-	-	-			1459
		gag Glu	-		-	-	-	-	-	-				-	_	1507

3

					Gly					Val					Lys	g tcg s Ser	1555
				Ile					Val					Asr		aac Asn	1603
								Asn					Glu			cag Gln	1651
I																gag Glu 455	1699
																cgc Arg	1747
													ttc Phe			aga Arg	1795
													gcg Ala 500				1843
													tcg Ser				1891
A													caa Gln				1939
													tgc Cys				1987
													cac His				2035
g A	ac sp	Cys	tca Ser 570	gag Glu	tgc Cys	ttc Phe	ccc Pro	ggc Gly 575	gtg Val	tca. Ser	gaa Glu	tct Ser	caa Gln 580	ccg Pro	gtc Val	gtc Val	2083

aga aag agg acg Arg Lys Arg Thr			His Leu Leu Gly
cgg gct ccc gag Arg Ala Pro Glu 600	• •		
Leu Asp Asp Cys	=	taa atgacttaaa	ccaggt atg gct gcc 2231 Met Ala Ala 625
gat ggt tat ctt of Asp Gly Tyr Leu 630	•		
cgc gag tgg tgg 6 Arg Glu Trp Trp 2 645			·
cag caa aag cag o Gln Gln Lys Gln 2 660			
tac ctc gga ccc f			
gcg gac gca gcg ( Ala Asp Ala Ala A 695			
aaa gcg ggt gac a Lys Ala Gly Asp 7 710	<del>-</del>		
ttt cag gag cgt c Phe Gln Glu Arg 3 725			
cga gca gtc ttc o Arg Ala Val Phe o 740			• • •
gtt gag gaa ggc ( Val Glu Glu Gly A			

cag to	-		-											2711
cag ca Gln Gl	-	-		-	_				-			-		2759
gag to Glu Se 80	r Val		_					 -			-			2807
gct gc Ala Al 820						_	_				-		-	2855
gca ga Ala As			-		-	-		 		-				2903
tgg ca Trp Hi	-	-												2951
acc cg Thr Ar			-	_	Pro				His					2999
atc to Ile Se	r Ser												_	3047
tac ag Tyr Se 900														3095
ttt to Phe Se														3143
cgg co														3191
gtc ac	-				_								-	3239

= =			ctt ccg tac gtc ctc Leu Pro Tyr Val Leu 975	3287
			ccg gcg gac gtg ttc Pro Ala Asp Val Phe 995	3335
Met Ile Pro Gln		* -	aat ggc agc caa gcc Asn Gly Ser Gln Ala 1010	3383
-	Ser Phe Tyr		ttc cct tct cag atg Phe Pro Ser Gln Met 1025	3431
	Asn Asn Phe	•	acc ttt gag gaa gtg Thr Phe Glu Glu Val 1040	3479
•		His Ser Gln Ser	ctg gac cgg ctg atg Leu Asp Arg Leu Met 1055	3527
•	Asp Gln Tyr		aac aga act caa aat Asn Arg Thr Gln Asn 1075	3575
Gln Ser Gly Ser	-		ttt agc cgt ggg tct Phe Ser Arg Gly Ser 1090	3623
	-		cta cct gga ccc tgt Leu Pro Gly Pro Cys 1105	3671
	Arg Val Ser		gac aac aac agc Asp Asn Asn Asn Ser 1120	3719
		Ser Lys Tyr Asn	ctc aat ggg cgt gaa Leu Asn Gly Arg Glu 1135	3767
		•	cac aaa gac gac gaa His Lys Asp Asp Glu 1155	3815

-	_	ttc Phe														3863
7.05	_,,			1160			,		1165			,	-	1170		
gcc	gga	gct	tca	aac	act	gca	ttg	gac	aat	gtc	atg	att	aca	gac	gaa	3911
Ala	Gly	Ala	Ser	Asn	Thr	Ala	Leu	Asp	Asn	Val	Met	Ile	Thr	Asp	Glu	
		J	175					1180				1	1185			
gag	gaa	att	aaa	gcc	act	aac	cct	gtg	gcc	acc	gaa	aga	ttt	ggg	acc	3959
Glu	Glu	Ile	Lys	Ala	Thr	Asn	Pro	Val	Ala	Thr	Glu	Arg	Phe	Gly	Thr	•
	1	190			•.	. 1	1195				1	200				
gtg	gca	gtc	aat	ttc	cag	agc	agc	agc	aca	gac	cct	gcg	acc	gga	gat	4007
Val	Ala	Val	Asn	Phe	Gln	Ser	ser	Ser	Thr	Asp	Pro	Ala	Thr	Gly	Asp	
1	1205				1	210				1	215					
ata	cat	gct	atq	gga	αca	tta	cct	aac	atα	ata	taa	caa	gat	aga	gac	4055
		Ala	-		_									-		
1220				_	225			3		230			<b>-</b>	-	235	
, ,		ctg	_					_						-		4103
Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	Pro	His	Thr	Asp	Gly	
			1	1240				]	245				1	1250		
cac	ttt	cac	ccg	tct	cct	ctt	atg	ggc	ggc	ttt	gga	ctc	aag	aac	ccg	. 4151
His	Phe	His	Pro	Ser	Pro	Leu	Met	Gly	Gly	Phe	Gly	Leu	Lys	Asn	Pro	
		1	255				1	1260		• .		1	.265			
cct	cct	cag	atc	ctc	atc	aaa	aac	acg	cct	gtt	cct	gcg	aat	cct	ccg	4199
Pro	Pro	Gln	Ile	Leu	Ile	Lys	Asn	Thr	Pro	Val	Pro	Ala	Asn	Pro	Pro	
	1	1270				1	275				1	.280				
aca	gag	ttt	tca	act	aca	aaq	ttt	act	tca	ttc	atc	acc	caa	tac-	tcc	4247
		Phe		-		_		-								
	1285					.290		•			.295			•		
		caa														4295
	_	Gln	Val			Glu	Ile	Glu			Leu	Gln	Lys	Glu	Asn	
1300	) .			1	.305					310				. ]	315	
agc	aag	cgc.	tgg	aat	ccc	gaa	gtg	cag	tac	aca	tcc	aat	tat	gca	aaa	4343
Ser	Lys	Arg	Trp	Asn	Pro	Glu	Val	Gln	Tyr	Thr	Ser	Asn	Tyr	Ala	Lys	
	-	-	-	320					325				_	1330		
tet	acc	aac	att	gat	ttt	act	ata	gac	aac	aat	gσa	ctt	tat	act	gag	4391
	_	Asn	_	_												
			335					340			1		345			

cct cgc ccc att ggc acc cgt tac ctt acc cgt ccc ctg taattacgtg 4440 Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu 1350 1355 ttaatcaata aaccggttga ttcgtttcag ttgaactttg gtctcctgtc cttcttatct 4500 tateggttae catggttata gettacacat taactgettg gttgegette gegataaaag 4560 acttacgtca tcgggttacc cctagtgatg gagttgccca ctccctctct gcgcgctcgc 4620 tegeteggtg gggeetgegg accaaaggte egcagaegge agagetetge tetgeeggee 4680 4718 ccaccgagcg agcgagcgcg cagagaggga gtgggcaa <210> 2 <211> 623 <212> PRT <213> AAV-1 <400> 2 Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 25 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 40 35 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 55 50 Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 75 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile

100

105

90

110

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly . 130 135 140

Asn 145	Lys	Val	Val	Asp	Glu 150	Cys	туr	Ile	Pro	155	Tyr	Leu	Leu	Pro	Lys 160
Thr	Gln	Pro	Glu	Leu 165	Gln	Trp	Ala	Trp	Thr 170	Asn	Met	Glu	Glu	туr 175	Ile
Ser	Ala	Суз	Leu 180	Asn	Leu	Ala	Glu	Arg 185	Lys	Arg	Leu	Val	Ala 190	Gln	His
Leu	Thr	His 195	Val	Ser	Gln		Gln 200	Glu	Gln	Asn	Lys	Glu 205	Asn	Leu	Asn
Pro	Asn 210	Ser	Asp	Ala	Pro	Val 215	Ile	Arg	Ser	Lys	Thr 220	Ser	Ala	Arg	Tyr
Met 225	Glu	Leu	Val	Gly	Trp 230	Leu	Val	Asp	Arg	Gly 235	Ile	Thr	Ser	Glu	Lys 240
				245	-				250				Asn	255	
			260					265			-		Ala 270	-	-
		275					280					285	Gly		
	290		-		-	295		-			300		Leu		
305	_	-			310	-		_		315			Gly	-	320
				325		ī			330				Gly	335	
		-	340	•				345	÷				Ala 350		
		355					360					365	Phe		
-	370	-				375					380		Met		•
ъуs 385	val.	vai⊥	GIU	SET	390	ъÀр	₩Ţ₫	тте	теп	395	grà	ser	Lys	AGT	400

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
405 410 415

- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445
- Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460
- Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480
- Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495
- Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510
- Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525
- Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 535 540
- Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560
- Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575
- Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590
- Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605
- Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 3

<211> 736

<212> PRT

<213> AAV-1

<400> 3

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 5 10 15

- Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30
- Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45
- Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60
- Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80
- Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95
- Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
  100 105 110
- Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125
- Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140
- Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly 145 150 155 160
- Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175
- Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190
- Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205
- Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220
- Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240
- Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu

245 250 255

Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270

- Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285
- His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300
- Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320
- Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335
- Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350
- Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365
- Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380
- Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400
- Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415
- Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430
- Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
  435 440 445
- Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 455 460
- Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480
- Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495
- Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn

500

505

510

Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys
515 520 525

Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540

Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560

Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg 565 570 575

Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590

Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln
595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 620

Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 630 635 640

Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655

Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 4

<211> 1872

<212> DNA

<213> AAV-1

<220> <221> CDS <222> (1)..(1869)

<400> 4 atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag 96 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu .25 20 aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 . 40 gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 55 qtc caa tqq cqc cqc qtq aqt aaq gcc ccg gag gcc ctc ttc ttt gtt Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 336 acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 120 125 115 ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag 480 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 150 155

act cag ccc gag ctg cag tgg gcg tgg act aac atg gag gag tat ata

Thr	Gln	Pro	Glu	Leu 165	Gln	Trp	Ala	Trp	Thr 170	Asn	Met	Glu	Glu	Туг 175	Ile	
-	-	-	ttg Leu 180		-	-		-						_		576
-			gtc Val	-	-		-		_		-			_	aac Asn	624
			gac Asp			-								-	tac. Tyr	672
_		_	gtc Val			_	-	-	-						-	720
_			cag Gln		-	_	-	-						-	•	768
		-	cgg Arg 260		_		-	-	-	-	-		-		_	816
	_		ctg Leu				_		_		_	-			-	864
-			gac Asp					-			-		•			912
			gaa Glu				_			-					-	960
_			ttc Phe		_	-					_			_	-	1008
	_		aag Lys 340				Ala	-	-		-		-			1056
ttc	tac	ggc	tgc	gtc	aac	tgg	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	1104

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 360 tgc gtc gac aag atg gtg atc tgg tgg gag gag ggc aag atg acg gcc 1152 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 370 375 aag gtc gtg gag tcc gcc aag gcc att ctc ggc ggc agc aag gtg cgc Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 390 gtg gac caa aag tgc aag tcg tcc gcc cag atc gac ccc acc ccc gtg 1248 . Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 atc gtc acc tcc aac acc aac atg tgc gcc gtg att gac ggg aac agc 1296 Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 430 acc acc ttc gag cac cag cag ccg ttg cag gac cgg atg ttc aaa ttt 1344 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 gaa ctc acc cgc cgt ctg gag cat gac ttt ggc aag gtg aca aag cag 1392 Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460 gaa gtc aaa gag ttc ttc cgc tgg gcg cag gat cac gtg acc gag gtg 1440 Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 gcg cat gag tto tac gtc aga aag ggt gga gcc aac aaa aqa ccc qcc Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 ccc gat gac gcg gat aaa agc gag ccc aag cgg gcc tgc ccc tca gtc 1536 Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 505 510 gcg gat cca tcg acg tca gac gcg gaa gga gct ccg gtg gac ttt gcc Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 gac agg tac caa aac aaa tgt tct cgt cac gcg ggc atg ctt cag atg 1632 Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 535 ctg ttt ccc tgc aag aca tgc gag aga atg aat cag aat ttc aac att 1680

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 550 555 tgc ttc acg cac ggg acg aga gac tgt tca gag tgc ttc ccc ggc gtg 1728 Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 575 565 tca gaa tct caa ccg gtc gtc aga aag agg acg tat cgg aaa ctc tgt Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 585 580 gcc att cat cat ctg ctg ggg cgg gct ccc gag att gct tgc tcg gcc 1824 Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 600 . 595 tgc gat ctg gtc aac gtg gac ctg gat gac tgt gtt tct gag caa taa 1872 Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 <210> 5 <211> 623 <212> PRT <213> AAV-1 <400> 5 Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 5. 10 1 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 40 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 55 50 Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 70 65 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190

Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 240

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255

Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270

Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285

Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300

Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315 320

Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro  $340 \hspace{1cm} 345 \hspace{1cm} 350$ 

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala

370 375 380

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400

- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
  435 440 445
- Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460
- Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480
- Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495
- Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510
- Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525
- Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 540
- Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 560
- Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575
- Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590
- Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605
- Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 6 <211> 1641 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1638) <400> 6 atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 1 10 15 gag cac etg ecg ggc att tet gac teg ttt gtg age tgg gtg gee gag Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 25 aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 40 45 gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt 240 Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 75 cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag 288 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95 acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg 384 Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 120 ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys

age gee tgt ttg aac etg gee gag ege aaa egg ete gtg geg eag eac 576 Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His ctg acc cac gtc agc cag acc cag gag cag aac aag gag aat ctg aac Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn ccc aat tot gac gog cot gto ato ogg toa aaa acc too gog ogc tac Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala tee aac teg egg tee eag ate aag gee get etg gae aat gee gge aag Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys atc atg gcg ctg acc aaa tcc gcg ccc gac tac ctg gta ggc ccc gct Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala ccg ccc gcg gac att aaa acc aac cgc atc tac cgc atc ctg gag ctg Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu aac ggc tac gaa cct gcc tac gcc ggc tcc gtc ttt ctc ggc tgg gcc Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 310 315 cag aaa agg ttc ggg aag cgc aac acc atc tgg ctg ttt ggg ccg gcc Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 acc acg ggc aag acc aac atc gcg gaa gcc atc gcc cac gcc gtg ccc Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro

22

340 345 350

ttc	tac	ggc	tgc	gto	aac	tgg	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	1104
															Asp	
	-	355				•	360					365			1.00	
							,					500				
tac	atc	asc	220	ato	ara	ato	taa	+~~	~~~	<b>~</b> ~ ~	~~~	224		~	gcc	1150
																1152
Cys		ASP	Lys	Met	vaı		Trp	Trp	GIU	GIu	-	ьys	Met	Thr	Ala	
	370			•		375					380					
			gag													1200
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg	
385					390					395					400	
gtg	gac	caa	aag	tgc	aag	tcg	tèc	gcc	cag	atc	gac	ccc	acc	ccc	gtg	1248
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Ala	Gln	Ile	Asp	Pro	Thr	Pro	Val	
				405					410		-			415		
atc	atc	acc	tcc	aac	acc	aac	at.g	tac	acc	ata	att	gac	aaa	aac	age	1296
			Ser													1270
110	, , ,		420		1111	ASIL	ric c		ATA	Val	116	Asp	_	ASII	ser	
							,	425					430			
															•	
			gag													1344
Thr	Thr		Glu	His	Gln	Gln		Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe	
		435					440					445				
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt,	ggc	aag	gtg	aca	aag	cag	1392
Glu	Leu	Thr	Arg	Arg	Leu	Glu	His	Asp	Phe	Gly	Lys	Val	Thr	Lys	Gln	
	450					455					460					
gaa	gtc	aaa	gag	ttc	ttc	cgc	tgg	gcg	caq	gat	cac	ata	acc	gag	ata	1440
			Glu													
465		•			470					475			•••-		480	
															400	
aca	cat	nan	ttc	tac	atc	202	220		~~~	~~~	330					1400
																1488
лта	nra	GIU	Phe		val	Arg	гàг	GIY		Ата	ASI	гÀг	Arg		Ala	
		•		485					490					495		
											•					
			gcg													1536
Pro	Asp	Asp	Ala	Asp	Lys	Ser	Glu	Pro	Lys	Arg	Ala	Cys	Pro	Ser	Val	
			<b>50</b> 0					505					510			
																÷
gcg	gat	cca	tcg	acg	tca	gac	gcg	gaa	gga	gct	ccq	gta	gac -	ttt	qcc	1584
			Ser													
	-	515					520		,			525	_			•
٠		•														
gac	agg	tat	ggc	tac	CG 3	taa	tto	tct	tee	202	++~	~~+	<b>~</b> ~~	~~-		1600
																1632
-up	A. y	- Y L	Gly	cys	vīd	тp	ьeu	ser	ser	Arg	ren	нта	Arg	GTÀ	GIN	

530

535

540

cct ctc tga
Pro Leu
545

1641 .

<210> 7 <211> 546

<212> PRT

<213> AAV-1

<400> 7

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp
1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

- Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220
- Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 240
- Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255
- Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270
- Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285
- Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300
- Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315 320
- Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335
- Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350
- Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 . 365
- Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 370 375 380
- Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400
- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
  420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

GIU	450	Thr	Arg	Arg	Leu	455	итэ	мар	·	GIY	460	vai	IIIL	пАЭ	GIII	
Glu 465	Val	Lys	Glu	Phe	Phe 470	Arg	Trp	Ala	Gln	Asp 475	His	Val	Thr	Glu	Val 480	
Ala	His	Glu	Phe	Tyr 485	Val	Arg	Lys	Gly	Gly 490	Ala	Asn	Lys	Arg	Pro 495	Ala	
Pro	Asp	Asp	Ala 500	Asp	Lys	Seŕ	Glu	Pro 505	Lys	Arg	Ala	Cys	Pro 510	Ser	Val	
Ala	Asp	Pro 515	Ser	Thr	Ser	Asp	Ala 520	Glu	Gly	Ala	Pro	Val 525	Asp	Phe	Ala	÷
Asp	Arg 530	Туr	Gly	Cys	Arg	Trp 535	Leu	Ser	Ser	Arg	Leu 540		Arg	Gly	Gln	•
Pro 545	Leu															
<212 <213 <220 <221	L> 12 2> DN 3> AA D> L> CI	AV-1 OS	(1197	7)												
<400	)> 8															
_		_	gtc Val													48
-		Ile	cag Gln 20	-	-	_	-	_						-	_	96
		-	cgg Arg		-		-	-	-	_	-		_		_	144
	_		ctg Leu					•	-		_	-				192

ccg Pro 65	ccc Pro	gcg Ala	gac Asp	att Ile	aaa Lys 70	acc Thr	aac Asn.	cgc Arg	atc Ile	tac Tyr 75	cgc Arg	atc Ile	ctg Leu	gag Glu	ctg Leu 80	240
			gaa Glụ													288
			ttc Phe 100													336
acc Thr	acg Thr	ggc Gly 115	aag Lys	acc Thr	aac Asn	atc Ile	gcg Ala 120	gaa Glu	gcc Ala	atc Ile	gcc Ala	cac His 125	gcc Ala	gtg Val	ccc Pro	384
			tgc Cys													432
			aag Lys													480
			gag Glu													528
			aag Lys 180													576
			tcc Ser													624
			gag Glu													672
			cgc Arg													720
			gag Glu													768

		-			-	•	-			-		aaa Lys	_		_	816
	-	-		_		•			_		-	tgc Cys 285			-	864
	-		_	-		-		_		-	-	gtg <b>V</b> al	-		_	912
•						-		_				atg Met		_	-	960
-			_	-		-		-	-			aat Asn				1008
-		-			_	-	-	_			-	ttc Phe			, ,	1056
	-			_	•	_	-	-		-		cgg Arg 365			_	1104
												gct Ala				1152
-	-	_	-			_	-	-	-	_	_	tct Ser			taa	1200

<210> 9 <211> 399

<212> PRT

<213> AAV-1

<400> 9

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 1 5 . 10 . 15

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala

											•				
Ser	Asn	Ser 35	Arg	Ser	Gln	Ile	Lys 40		Ala	Leu	Asp	Asn 45	Ala	Gly	Ly
Ile	Met 50	Ala	Leu	Thr	Lys	Ser 55	Ala	Pro	Asp	Туr	Leu 60	Val	Gly	Pro	Al
Pro 65	Pro	Ala	Asp	Ile	Lys 70	Thr	Asn	Arg	Ile	Туг 75	Arg	Ile	Leu	Glu	Le 8
Asn	Gly	Tyr	Glu	Pro 85	Ala	Tyr	Ala	Gly	Ser 90	Val	Phe	Leu	Gly	Trp 95	Al
Gln	Lys	Arg	Phe 100	Gly	Lys	Arg	Asn	Thr 105	Ile	Trp	Leu	Phe	Gly 110	Pro	Al
Thr	Thr	Gly 115	Lys	Thr	Asn	Ile	Ala 120	Glu	Ala	Ile	Ala	His 125	Ala	Val	Pr
Phe	туг 130	Gly	Cys	Val	Asn	1 rp	Thr	Asn	Glu	Asn	Phe 140	Pro	Phe	Asn	Asj
Cys 145	Val	Asp	Lys	Met	Val 150	Ile	Trp	Trp	Glu	Glu 155	Gly	Lys	Met	Thr	Ala 16
Lys	Val	Val	Glu	Ser 165	Ala	Lys	Ala	Île	Leu 170	Gly	Gly	Ser	Lys	Val 175	Ar
Val	Asp	Gln	Lys 180	Cys	Lys	Ser	Ser	Ala 185	Gln	Ile	Asp	Pro	Thr 190	Pro	Va.
lle	Val	Thr 195	Ser	Asn	Thr	Asn	Met 200	Cys	Ala	Val	Ile	Asp 205	Gly	Asn	Se
Thr	Thr 210	Phe	Glu	His	Gln	Gln 215	Pro	Leu	Gln	Asp	Arg 220	Met	Phe	Lys	Phe
Glu 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys	Val	Thr	Lys	Gl: 240
Glu	Val	Lys	Glu	Phe 245	Phe	Arg	Trp	Ala	Gln 250	Asp	His	Val	Thr	Ġlu 255	Va:
Ala	His		Phe 260	Tyr	Val	Arg	Lys	Gly 265	Gly	Ala	Asn	Lys	Arg 270	Pro	Ala

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val

275 280 285

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 300

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 305 310 315 320

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 325 330 335

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 340 345 350

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 355 360 365

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 370 375 380

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 385 390 395

<210> 10

<211> 969

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1)..(966)

<220>

<221> misc\_feature

<222> (943)..(944)

<223> minor splice site

<400> 10

atg gag ctg gtc ggg tgg ctg gtc ggc atc acc tcc gag aag. 48
Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys

1 5 10 15

cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct 96 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 20 25 30

tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 144

Ser	Asn	Ser 35	Arg	Ser	Gln	Ile	Lys 4 <sub>.</sub> 0	Ala	Ala	Leu	Asp	Asn 45	Ala	Gly	Lys	
	_		_	acc Thr					-		_	•			-	192
_			_	att Ile							-		_		_	240
			-	cct Pro 85	-		-			_					•	288
-				ggg Gly	_	-					•			_	•	336
	-		-	acc Thr				-	-		-		-			384
			-	gtc Val											•	432
				atg Met												480
_				tcc Ser 165	-	_	_					-	_		•	528
	-		_	tgc Cys	-	-		-	-		•					576
				aac Asn												624
				cac His	-	-	-	-	-	-		_				672
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt	ggc	aag	gtg	aca	aag	cag	720

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln . 230 235 225 gaa gtc aaa gag ttc ttc cgc tgg gcg cag gat cac gtg acc gag gtg 768 Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 245 gcg cat gag ttc tac gtc aga aag ggt gga gcc aac aaa aga ccc gcc 816 Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 260 265 ccc gat gac gcg gat aaa agc gag ccc aag cgg gcc tgc ccc tca gtc Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 280 gcg gat cca tcg acg tca gac gcg gaa gga gct ccg gtg gac ttt gcc 912 Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 290 295 300 gac agg tat ggc tgc cga tgg tta tct tcc aga ttg gct cgá gga caa 960 Asp Arg Tyr Gly Cys Arg Trp Leu Ser Ser Arg Leu Ala Arg Gly Gln 310 315 cct ctc tga 969 Pro Leu <210> 11 <211> 322 <212> PRT <213> AAV-1 Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 50 Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 75

Asn	.Gly	Tyr	Glu	Pro 85	Ala	Туr	Ala	Gly	Ser 90		Phe	Leu	Gly	Trp 95	Ala
Gln	Lys	Arg	Phe 100	Gly	Lys	Arg	Asn	Thr 105	Ile	Trp	Leu	Phe	Gly 110	Pro	Ala
Thr	Thr	Gly 115	Lys	Thr	Asn	Ile	Ala 120	Glu	Ala	Ile	Ala	His 125	Ala	Val	Pro
Phe	Туг 130	Gly	Cys	Val	Asn	Trp 135	Thr	Asn	Glu	Asn	Phe 140	Pro	Phe	Asn	Asp
Cys 145	Val	Asp	Lys	Met	Val 150	Ile	Тrр	Trp	Glu	Glu 155	Gly	Lys	Met	Thr	Ala 160
Lys	Val	Val	Glu	Ser 165	Ala	Lys	Ala	Ile	Leu 170	Gly	Gly	Ser	Lys	Val 175	Arg
Val	Asp	Gln	Lys 180	Cys	Lys	ser	Ser	Ala 185	Gln	Ile	Asp	Pro	Thr 190	Pro	Val
Ile	Val	Thr 195	Ser	Asn	Thr	Asn	Met 200	Cys	Ala	Val	Ile	Asp 205	Gly	Asn	Ser
Thr	Thr 210	Phe	Glu	His	Gln	Gln 215	Pro	Leu	Gln	Asp	Arg 220	Met	Phe	Lys	Phe
Glu 225	Leu	Thr	Arg	Arġ	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys ·	Val	Thr	Lys	Gln 240
Glu	Val	Lys	Glu	Phe 245	Phe	Arg	Trp	Ala	Gln 250	Asp	His	Val	Thr	Glu 255	Val
Ala	His	Glu	Phe 260	Tyr	Val ,	Arg	Lys	Gly 265	Gly	Ala	Asn	Lys	Arg 270	Pro	Ala
	Asp	275	•				280		-			285			
Ala	Asp 290	Pro	Ser	Thr	Ser	Asp 295	Ala	Glu	Gly	Ala	Pro 300	Val	Asp	Phe	Ala
Asp 305	Arg	Tyr	Gly	Cys	Arg 310	Trp	Leu	Ser	Ser	Arg 315	Leu	Ala	Arg	Gly	Gln 320
Pro	Leu														

<210> 12 <211> 2211 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(2208) <400> 12 atg gct gcc gat ggt tat ctt cca gat tgg ctc gag gac aac ctc tct. 48 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser 10 gag ggc att cgc gag tgg tgg gac ttg aaa cct gga gcc ccg aag ccc Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 aaa gcc aac cag caa aag cag gac gac ggc cgg ggt ctg gtg ctt cct Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 ggc tac aag tac ctc gga ccc ttc aac gga ctc gac aag ggg gag ccc 192 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 55 gtc aac gcg gcg gac gca gcg gcc ctc gag cac gac aag gcc tac gac Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 cag cag ctc aaa gcg ggt gac aat ccg tac ctg cgg tat aac cac gcc 288 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala gac gcc gag ttt cag gag cgt ctg caa gaa gat acg tct ttt ggg ggc Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly 100 105 aac ctc ggg cga gca gtc ttc cag gcc aag aag cgg gtt ctc gaa cct 384 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125 ctc ggt ctg gtt gag gaa ggc gct aag acg gct cct gga aag aaa cgt Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 ccg gta gag cag tcg cca caa gag cca gac tcc tcc tcg ggc atc ggc

Pro 145	Val	Glu	Gln	Ser	Pro 150	Gln	Glu	Pro	Asp	Ser 155	Ser	Ser	Gly	Ile	Gly 160	
-			-	_	ccc Pro	-		_	-					_		528
	-				gtc Val								-			576 ·
-			-	-	gtg Val					_	-					624
_					aat Asn										gcc. Ala	672
					tgc Cys 230	_				_		-	•	-		720
		-		-	acc Thr		-	-								768
	_				agt Ser	_		-		-	_		-	Asn		816
				-	acc Thr						-			-		864
					cca Pro											912
					aag Lys 310											960
-	_		1	_	acg Thr		_		_				-			1008
ctt	acc	agc	acg	gtt	caa	gtc	ttc	tcg	gac	tcg	gag	tac	cag	ctt	ccg	1056

Leu	Thr	Ser	Thr 340	Val	Gln	Val	Phe	ser 345	Asp	Ser	Glu	Туг	Gln 350	Leu	Pro	
		ctc Leu 355														1104
		ttc Phe														1152
		gcc Ala				Ser										1200
		atg Met														1248
		gtg Val														1296
		atg Met 435														1344
		aat Asn														1392
		tct Ser														1440
		tgt Cys														1488
		agc Ser														1536
		gaa Glu 515						Gly								1584
gac	gac	gaa	gac	aag	ttc	ttt	ccc	atg	agc	ggt	gtc	atg	att	ttt	gga	1632

	-																
Į	Asp	Asp 530	Glu	Asp	Lys ·	Phe	Phe 535	Pro	Met	Ser	Gly	Val 540	Met	Ile	Phe	Gly	
I			-												atg Met		1680
		-	-		-									Thr	gaa Glu 575		1728
															cct Pro		1776
															tgg Trp		1824
		-	-												cct Pro		1872
3		-													gga Gly		1920
	_			Pro											cct Pro 655		1968
															atc Ile		2016
															ctg Leu		2064
		-													tcc Ser		2112
7		-			_		-	-				_			gga Gly		2160
1	at	act	gag	cct	cgc	ccc	att	ggc	acc	cgt	tac	ctt	acc	cgt	ccc	ctg	2208

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

taa 2211

<210> 13

<211> 736

<212> PRT

<213> AAV-1

<400> 13

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser 1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly 100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 · 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly 145 150 155 160

Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175

Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190

Ala	Thr	Pro 195	Ala	Ala	Val	Gly	Pro 200	Thr	Thr	Met	Ala	ser 205	Gly	Gly	Gly
Ala	Pro 210	Met	Ala	Asp	Asn	Asn 215	Glu	Gly	Ala	Asp	Gly 220	Val	Gly	Asn	Ala
Ser 225	Gly	Asn	Trp	His	Суз 230	Asp	Ser	Thr	Trp	Leu 235	Gly	Asp	Arg	Val	Ile 240
Thr	Thr	Ser	Thr	Arg 245	Thr	Trp	Ala	Leu	Pro 250	Thr	Туr	Asn	Asn	His 255	Leu
Tyr	Lys	Gln	11e 260	Ser	Ser	Ala	Ser	Thr 265	G] À	Ala	Ser	Asn	Asp 270	Asn	His
Tyr	Phe	Gly 275	Tyr	Ser	Thr	Pro	Trp 280	Gly	Туг	Phe	Asp	Phe 285	Asn	Arg	Phe
His	Cys 290	His	Phe	Ser	Pro	Arg 295	Asp	Trp	Gln	Arg	Leu 300	Ile	Asn	Asn	Asn
Trp 305	Gly	Phe	Arg	Pro	Lys 310	Arg	Leu	Asn	Phe	Lys 315	Leu	Phe	Asn	Ile	Gln 320
Val	Lys	Glu	Val	Thr 325	Thr	Asn	Asp	Gly	Val 330	Thr	Thr	Ile	Ala	Asn 335	Asn
Leu	Thr	Ser	Thr 340	Val	Gln	Val	Phe	ser 345	Asp	ser	Glu	Туr	Gln 350	Leu	Pro
Tyr	Val	Leu 355	Gly	Ser	Ala	His	Gln 360	Gly	Cys	Leu	Pro	Pro 365	Phe	Pro	Ala
Asp	·Val 370	Phe	Met	Ile	Pro	Gln 375	Tyr	Gly	Туг	Leu	Thr 380	Leu	Asn	Àsn	Gly
Ser 385	Gln	Ala <sup>.</sup>	Val	Gly	Arg 390	Ser	Ser	Phe	туг	Cys 395	Leu	Glu	Tyr	Phe.	Pro 400
Ser	Gln	Met	Leu	Arg 405	Thr	Gly	Asn	Asn	Phe 410	Thr	Phe	Ser	Тyr	Thr 415	Phe
Glu	Glu	Val	Pro 420	Phe	His	Ser	Ser	Туг 425	Ala	His	Ser	Gln	ser 430	Leu	Asp
Arg	Leu	Met 435	Asn	Pro	Leu	Ile	Asp 440	Gln	туг	Leu	Tyr	Tyr 445	Leu	Asn	Arg

Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 . 455 460

- Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480
- Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495
- Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn 500 505 510
- Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525
- Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540
- Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 560
- Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg
  565 570 575
- Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590
- Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln
  595 600 605
- Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 620
- Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 630 630 640
- Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655
- Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670
- Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685
- Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu 725 730 735

<210> 14

<211> 1800

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1) .. (1797)

<400> 14

acg gct cct gga aag aaa cgt ccg gta gag cag tcg cca caa gag cca 48
Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro
1 5 10 15

gac tcc tcc tcg ggc atc ggc aag aca ggc cag cag ccc gct aaa aag 96
Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys
20 25 30

aga ctc aat ttt ggt cag act ggc gac tca gag tca gtc ccc gat cca 144
Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro
35 40 45

caa cct ctc gga gaa cct cca gca acc ccc gct gct gtg gga cct act 192 Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr 50 55 60

aca atg gct tca ggc ggt ggc gca cca atg gca gac aat aac gaa ggc 240 Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 75 80

gcc gac gga gtg ggt aat gcc tca gga aat tgg cat tgc gat tcc aca 288
Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr
85 90 95

tgg ctg ggc gac aga gtc atc acc acc agc acc cgc acc tgg gcc ttg 336
Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu
100 105 110

41

		agc Ser						Thr		432
		gat Asp								480
		ctc Leu								528
		ctc Leu								576
		acc Thr 195								624
-	-	gag Glu								672
		cct Pro								720
		acg Thr								768
	_	ctg Leu	-							816
		ttc Phe 275	-			-				864
		agc Ser								912
	Leu	tat Tyr								960

	-	gac Asp	-	•		•	-				-		-		_	1008
,		aaa Lys							Cys			-	_	-	-	1056
		aca Thr 355			-											1104
-		aaa Lys														1152
	_	atg Met	-				•	-	-	-	-				-	1200
•		gtc Val	_						-	-						1248
-	•	gac Asp		-	-			-	-		-			-		1296
		gtg Val 435	-		-	Arg					-	-				1344
•	-	agc Ser		-					-			-	_		-	1392
		ggc Gly			-			-,	-			_	_			1440
		gcc Ala											_			1488
	-	ggc Gly					_	-	-			•				1536

	aac Asn	-		-							-			-		1584
-	ttt Phe 530	•												-		1632
-	att Ile				-	_		-								1680
-	gtg Val	_									-		-	-		1728
	gtg Val	_									_					1776
-	tac Tyr			-		-	taa									1800
		333														
<21:	0> 15 1> 59 2> PF 3> AF	5 99 RT										·				
<213 <213 <213	1> 59 2> PF 3> AF	5 99 RT AV-1														
<213 <213 <213	1> 59 2> PF	5 99 RT AV-1	Gly	Lys 5	Lys	Arg	Pro	Val	Glu 10	Gln	Ser	Pro	Gln	Glu 15	Pro	
<213 <213 <213 <400 Thr	1> 59 2> PF 3> AF 0> 15	5 99 RT NV-1 5 Pro	-	· 5	-				10 Gly					15		
<21: <21: <21: <400 Thr 1	1> 59 2> PF 3> AA 0> 19 Ala	Ser	Ser 20	Gly	Ile	Gly	Lys	Thr 25	10 Gly	Gln	Gln	Pro	Ala 30	15 Lys	Lys	
<21: <21: <21: <40: Thr 1 Asp	1> 59 2> PF 3> AF 3> 19 Ala Ser	5 99 RT W-1 5 Pro Ser Asn 35	Ser 20 Phe	Gly	Ile Gln	Gly	Lys Gly 40	Thr 25 Asp	10 Gly Ser	Gln Glu	Gln Ser	Pro Val 45	Ala 30 Pro	15 Lys Asp	Lys Pro	
<21: <21: <400 Thr 1 Asp Arg	1> 59 2> PF 3> AA 0> 15 Ala Ser Leu Pro	Ser Asn 35	Ser 20 Phe Gly	Gly Gly Glu	Ile Gln Pro	Gly Thr Pro 55	Lys Gly 40 Ala	Thr 25 Asp	10 Gly Ser Pro	Gln Glu Ala	Gln Ser Ala 60	Pro Val 45 Val	Ala 30 Pro Gly	15 Lys Asp	Lys Pro Thr	

85 90 95

Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu 100 105 110

- Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr 115 120 125
- Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
  130 135 140
- Tyr Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp 145 150 155 160
- Gln Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn 165 170 175
- Phe Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly 180 185 190
- Val Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser 195 200 205
- Asp Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly 210 215 220
- Cys Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly 225 230 235 240
- Tyr Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe
  245 250 255
- Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn 260 265 270
- Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr 275 280 285
- Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln 290 295 300
- Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln 305 310 315 320
- Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val 325 330 335
- Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val

340 345 350

Ser Lys Thr Lys Thr Asp Asn Asn Ser Asn Phe Thr Trp Thr Gly 355 360 365

Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly 370 375 380

Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met 385 390 395 400

Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr 405 410 415

Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr 420 425 430

Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln 435 440 445

Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala 450 455 460

Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro 465 470 475 480

Ilé Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro 485 490 495

Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile 500 505 510

Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 515 520 525

Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 530 535 540

Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 545 550 555 560

Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe 565 570 575

Thr Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr 580 585 590

Arg Tyr Leu Thr Arg Pro Leu

<21	0> 1	6														
<21	1> 1	605														
<21	2> D	NA														
<21	3> A	AV-1														
<22	მ>															
<22	1> C	DS												•		
<22	2> (	1)	(160	2)												
<40	0> 1	6														
atg	gct	tca	ggc	ggt	ggc	gca	сса	atg	gca	gac	aat	aac	gaa	ggc	gcc	48
Met	Ala	Ser	Gly	Gly	Gly	Ala	Pro	Met	Ala	Asp	Asn	Asn	Glu	Gly	Ala	
1				5					10					15		
gac	gga	gtg	ggt	aaț	gcc	tca	gga	aat	tgg	cat	tgc	gat	tcc	aca	tgg	96
Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Thr	Trp	
٠.			20					25					30			
ctg	ggc	gac	aga	gtc	atc	acc	acc	agc	acc	cgc	acc	tgg	gcc	ttg	ccc	144
					Ile								-	_		
		35					40		÷			45				
acc	tac	aat	aac	cac	ctc	tac	aag	caa	atc	tcc	agt	gct	tca	acg	ggg	192
Thr	Tyr	Asn	Asn	His	Leu	Tyr	Lys	Gln	Ile	Ser	Ser	Ala	Ser	Thr	Gly	
	50					55					60					
gcc	agc	aac	gac	aac	cac	tac	ttc	ggc	tac	agc	acc	ccc	tgg	ggg	tat	240
Ala	Ser	Asn	Asp	Asn	His	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	
65					70					75	•				80	
ttt	gat	ttc	aac	aga	ttc	cac	tgc	cac	ttt	tca	сса	cgt	gac	tgg	cag	288
Phe	Asp	Phe	Asn	Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	
				85					90					95		
cga	ctc	atc	aac	aac	aat	tgg	gga	ttc	cgg	ccc	aag	aga	ctc	aac	ttc	336
Arg	Leu	Ile	Asn	Asn	Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	
		•	100					105					110			
aaa	ctc	ttc	aac	atc	caa	gtc	aag	gag	gtc	acg	acg	aat	gat	ggc	gtc	384
					Gln											
		115	÷				120				•	125	-	-		
aca	acc	atc	gct	aat	aac	ctt	acc	agc	acq	att	caa	atc	ttc	tca	gac	432
					Asn											

		tac Tyr									480
		ccg Pro									528
	_	ctc Leu									576
-	_	gaa Glu 195									624
		agc Ser									672
	_	cag Gln								tac Tyr 240	720
		tac Tyr									768
-	-	ttg Leu	-	-							816
		aac Asn 275									864
aaa Lys	aca Thr 290	aaa Lys	aca Thr				aat Asn				912
		tat Tyr									960
		gcc Ala				,					1008

	-	-						_	-		-		aac Asn 350		_	1056
•	_		_	_			-	-		-			gcc Ala			1104
		-		-	-					-	-		ttc Phe	_	-	1152
-			-					- •			_	_	gga Gly	-		1200
		_				_	-	-			_	_	ggt Gly			1248
							-					-	tct Ser 430			1296
_						-		-			-		ctc Leu			1344
	_		_		-			_					gct Ala		_	1392
	-											-	agt Ser		-	1440
	_			-	_		_		_	_	-		aat Asn		-	1488
	_		Thr				_			-		-	gat Asp 510			1536
								•					ggc Gly		-	1584

tac ctt acc cgt ccc ctg taa Tyr Leu Thr Arg Pro Leu 530 1605

<210> 17 <211> 534 <212> PRT <213> AAV-1

<400> 17

Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala 1 5 10 15

Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp 20 25 30

Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro  $35 \hspace{1cm} 40 \hspace{1cm} 45$ 

Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly 50 55 60

Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr 65 70 75 80

Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln 85 90 95

Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe 100 105 110

Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val 115 : 120 125

Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp 130 135 140

Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys 145 150 155 160

Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr 165 170 175

Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr 180 185 190

Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met Ser' Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys 

Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys 450 455 460

Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu 465 470 475 480

Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu
485 490 495

Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr 500 505 510

Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg 515 520 525

Tyr Leu Thr Arg Pro Leu 530

<210> 18

<211> 4681

<212> DNA

<213> aav-2

<400> 18

tcacaaagac cagaaatggc gccggaggcg ggaacaaggt ggtggatgag tgctacatcc 780 ccaattactt gctccccaaa acccagcctg agctccagtg ggcgtggact aatatggaac 840 agtatttaag cgcctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900 cgcacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat tctgatgcgc 960 cggtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca 1020 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatac atctccttca 1080 atgcggcctc caactcgcgg tcccaaatca aggctgcctt ggacaatgcg ggaaagatta 1140 tgagcctgac taaaaccgcc cccgactacc tggtgggcca gcagcccgtg gaggacattt 1200 ccagcaatcg gatttataaa attttggaac taaacqqqta cqatccccaa tatqcqqctt 1260 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320 ggcctgcaac taccgggaag accaacateg cggaggecat ageceacaet gtgcccttet 1380 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440 tgatctggtg ggaggagggg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500 teggaggaag caaggtgege gtggaceaga aatgeaagte eteggeeeag atagaeeega 1560 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620 ccttcgaaca ccagcagccg ttgcaagacc ggatgttcaa atttqaactc acccqccqtc 1680 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cggtgggcaa 1740 aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaagggtgga gccaagaaaa 1800 gacccgcccc cagtgacgca gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc 1860 agccatcgac gtcagacgcg gaagcttcga tcaactacgc agacaggtac caaaacaaat 1920 gttctcgtca cgtgggcatg aatctgatgc tgtttccctg cagacaatgc gagagaatga 1980 atcagaattc aaatatctgc ttcactcacg gacagaaaga ctgtttagag tgctttcccg 2040 tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgta tcagaaactg tgctacattc 2100 atcatateat gggaaaggtg ceagaegett geaetgeetg egatetggte aatgtggatt 2160

tggatgactg catctttgaa caataaatga tttaaatcag gtatggctgc cgatggttat 2220 cttccagatt ggctcgagga cactctctct gaaggaataa gacagtggtg gaagctcaaa 2280 cctggcccac caccaccaaa gcccgcagag cggcataagg acgacagcag gggtcttgtg 2340 cttcctgggt acaagtacct cggaccttc aacggactcg acaagggaga gccggtcaac 2400 gaggcagacg ccgcggccct cgagcacgac aaagcctacg accggcagct cgacagcgga 2460 gacaacccgt acctcaagta caaccacgcc gacgcggagt ttcaggagcg ccttaaagaa 2520 qatacqtctt ttqqqqqcaa cctcqqacqa gcaqtcttcc aggcqaaaaa qaqqqttctt 2580 gaacctctcg gcctggttga ggaacctgtt aagacggctc cgggaaaaaa gaggccggta 2640 gagcactete etgtggagee agacteetee tegggaaceg gaaageeggg ecagcageet 2700 qcaagaaaaa gattgaattt tggtcagact ggagacgcag actcagtacc tgacccccag 2760 cctctcggac agccaccage agcccctct ggtctgggaa ctaatacgat ggctacaggc 2820 aqtqqcqcac caatgqcaga caataacgag ggcqccqacg gagtqgqtaa ttcctccgga 2880 aattggcatt gcgattccac atggatgggc gacagagtca tcaccaccag cacccgaacc 2940 tgggccctgc ccacctacaa caaccacctc tacaaacaaa tttccagcca atcaggagcc 3000 tegaaegaca ateaetaett tygetaeage aeeeettygg gytattttga etteaaeaga 3060 ttccactgcc actttccacc acgtgactgg caaagactca tcaacaacaa ctggggattc 3120 cgacccaaga gactcaactt caacctcttt aacattcaag tcaaagaggt cacgcagaat 3180 gacggtacga cgacgattgc caataacctt accagcacgg ttcaggtgtt tactgactcg 3240 gagtaccage tecegtacgt ceteggeteg gegeateaag gatgeeteee geegtteeca 3300 gcagacgtct tcatggtgcc acagtatgga tacctcaccc tgaacaacgg gagtcaggca 3360 gtaggacget etteatttta etgeetggag taettteett eteagatget gegtaeegga 3420 aacaacttta cottoagota cacttttgag gacgttoott tocacagoag ctacgotoac 3480 agccagagtc tggaccgtct catgaatect cteategace agtacetgta ttaettgage 3540 agaacaaaca ctccaagtgg aaccaccacg cagtcaaggc ttcagttttc tcaggcccca 3600

gccagtgaca ttcgggacca gtctaggaac tggcttcctg gaccctgtta ccgccagcag 3660 cgagtatgaa agacatctgc ggataacaac aacagtgaat actcgtggac tggagctacc 3720 aagtaccacc tcaatggcag agactetetg gtgaateegg ggccegecat ggcaageeac 3780 aaggacgatg aagaaaagtt ttttcctcag agcggggttc tcatctttgg gaagcaaggc 3840 tcagagaaaa caaatgtgaa cattgaaaag gtcatgatta cagacgaaga ggaaatccca 3900 acaaccaatc ccgtggctac ggagcagtat ggttctgtat ctaccaacct ccagagaggc 3960 aacagacaag cagctaccgc agatgtcaac acacaaggcg ttcttccagg catggtctgg 4020 caggacagag atgtgtacct tcaggggccc atctgggcaa agattccaca cacggacgga 4080 cattttcacc cctctcccct catgggtgga ttcggactta aacaccctcc tccacagatt 4140 ctcatcaaga acaccccggt acctgcgaat ccttcgacca ccttcagtgc ggcaaagttt 4200 getteettea teacacagta etceaeggga caeggteage gtggagateg agtgggaget 4260 gcagaacgaa aacagcaaac gctggaatcc cgaaattcag tacacttcca actacaacaa 4320 gtctgttaat cgtggacttt accgtggata ctaatggcgt gtattcagag cctcgcccca 4380 ttggcaccag atacctgact cgtaatctgt aattgcttgt taatcaataa accgtttaat 4440 togtttcagt tgaactttgg tototgcgta tttctttctt atctagtttc catggctacg 4500 tagataagta gcatggcggg ttaatcatta actacaagga acccctagtg atggagttgg 4560 ccactccctc tctgcgcgct cgctcgctca ctgaggccgg gcgaccaaag gtcgcccgac 4620 gcccgggctt tgccccggcg gcctcagtga gcgagcgagc gcgcaqaqaq qqaqtqqqca 4680 4681

<210> 19

<211> 4683

<212> DNA

<213> aav-6

<400> 19

ttggccacte cetetetgeg egetegeteg etcactgagg eegggegace aaaggtegee 60

cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgcag agagggagtg 120 gccaactcca tcactagggg ttcctggagg ggtggagtcg tgacgtgaat tacgtcatag 180 ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacaccat 240 gtggtcacgc tgggtattta agcccgagtg agcacgcagg gtctccattt tgaagcggga 300 ggtttgaacg cgcagcgcca tgccggggtt ttacgagatt gtgattaagg tccccagcga 360 ccttgacgag catctgcccg gcatttctga cagctttgtg aactgggtgg ccgagaagga 420 atgggagttg ccgccagatt ctgacatgga tctgaatctg attgagcagg cacccctqac 480 cgtggccgag aagctgcagc gcgacttcct ggtccactgg cgccgcgtga gtaaggcccc 540 ggaggccctc ttctttgttc agttcgagaa gggcgagtcc tacttccacc tccatattct 600 ggtggagace acgggggtca aatccatggt gctgggccgc ttcctgagtc agattagcga 660 caagetggtg cagaccatet accgegggat egageegace etgeecaact ggttegeggt 720 gaccaagacg cgtaatggcg ccggaggggg gaacaaggtg gtggacgagt gctacatccc 780 caactacete etgeccaaga eteageeega getgeagtgg gegtggaeta acatggagga 840 gtatataagc gcgtgtttaa acctggccga gcgcaaacgg ctcgtggcgc acgacctgac 900 ccacgtcagc cagacccagg agcagaacaa ggagaatctg aaccccaatt ctgacgcgcc 960 tgtcatccgg tcaaaaacct ccgcacgcta catggagctg gtcgggtggc tggtggaccg 1020 gggcatcacc teegagaage agtggateea ggaggaeeag geetegtaca teteetteaa 1080 egeogeotec aactegeggt eccagateaa ggeogetetg gacaatgeeg geaagateat 1140 ggcgctgacc aaatccgcgc ccgactacct ggtaggcccc gctccgcccg ccgacattaa 1200 aaccaaccgc atttaccgca teetggaget gaacggetae gaecetgeet acgeeggete 1260 cgtctttctc ggctgggccc agaaaaggtt cggaaaacgc aacaccatct ggctgtttgg 1320 gccggccacc acgggcaaga ccaacatcgc ggaagccatc gcccacgccg tgcccttcta 1380 eggetgegte aactggacca atgagaactt teeetteaac gattgegteg acaagatggt 1440 gatctggtgg gaggagggca agatgacggc caaggtcgtg gagtccgcca aggccattct 1500

eggeggeage aaggtgegeg tygaceaaaa gtgeaagteg teegeeeaga tegateeeac 1560 eccegtgate gteaceteca acaceaacat gtgegeegtg attgaeggga acageaceae 1620 cttcgagcac cagcagccgt tgcaggaccg gatgttcaaa tttgaactca cccgccgtct 1680 ggagcatgac tttggcaagg tgacaaagca ggaagtcaaa gagttettee getgggegea 1740 ggatcacgtg accgaggtgg cgcatgagtt ctacgtcaga aagggtggag ccaacaacag 1800 accegeeece gatgaegegg ataaaagega geeeaagegg geetgeeeet eagtegegga 1860 tocatogacg toagacgogg aaggagotoc ggtggacttt gccgacaggt accaaaacaa 1920 atgttctcgt cacgcgggca tgcttcagat gctgtttccc tgcaaaacat gcgagagaat 1980 gaatcagaat ttcaacattt gcttcacgca cgggaccaga gactgttcag aatgtttccc 2040 cggcgtgtca gaatctcaac cggtcgtcag aaagaggacg tatcggaaac tctgtgccat 2100 teateatetg etggggeggg etecegagat tgettgeteg geetgegate tggteaacgt 2160 ggatctggat gactgtgttt ctgagcaata aatgacttaa accaggtatg gctgccgatg 2220 gttatcttcc agattggctc gaggacaacc tctctgaggg cattcggcag tggtgggact 2280 tgaaacctgg agccccgaaa cccaaagcca accagcaaaa gcaggacgac ggccggggtc 2340 tggtgcttcc tggctacaag tacctcggac ccttcaacgg actcgacaag ggggagcccg 2400 tcaacgcggc ggatgcagcg gccctcgagc acgacaaggc ctacgaccag cagctcaaag 2460 egggtgacaa teegtacetg eggtataaee aegeegaege egagttteag gagegtetge 2520 aagaagatac gtottttggg ggcaacotog ggogagcagt ottocaggoo aagaagaggg 2580 ttetegaace tittggtetg gitgaggaag gigetaagae ggeteetgga aagaaacgte 2640 eggtagagea gtegecaeaa gagecagaet eeteeteggg cattggeaag acaggecage 2700 agcccgctaa aaagagactc aattttggtc agactggcga ctcagagtca gtccccgacc 2760 cacaacctct cggagaacct ccagcaaccc ccgctgctgt gggacctact acaatggctt 2820 caggoggtgg cgcaccaatg gcagacaata acgaaggcgc cgacggagtg ggtaatgcct 2880 caggaaattg gcattgcgat tccacatggc tgggcgacag agtcatcacc accagcaccc 2940

gaacatgggc cttgcccacc tataacaacc acctctacaa gcaaatctcc agtgcttcaa 3000 cgggggccag caacgacaac cactacttcg gctacagcac cccctggggg tattttgatt 3060 tcaacagatt ccactgccat ttctcaccac gtgactggca gcgactcatc aacaacaatt 3120 ggggattccg gcccaagaga ctcaacttca agctcttcaa catccaagtc aaggaggtca 3180 cgacgaatga tggcgtcacg accatcgcta ataaccttac cagcacggtt caagtcttgt 3240 cggactcgga gtaccagttc ccgtacgtcc tcggctctgc gcaccagggc tgcctccctc 3300 cgttcccggc ggacgtgttc atgattccgc agtacggcta cctaacgctc aacaatggca 3360 gccaggcagt gggacgctca tccttttact gcctggaata tttcccatcg cagatgctga 3420 gaacgggcaa taactttacc ttcagctaca ccttcgagga cgtgcctttc cacagcagct 3480 acgcgcacag ccagagcctg gaccggctga tgaatcctct catcgaccag tacctgtatt 3540 acctgaacag aactcacaat cagtccggaa gtgcccaaaa caaggacttg ctgtttagcc 3600 gtgggtetec agetggeatg tetgtteage ceaaaaaetg getaeetgga eeetgttaee 3660 ggcagcagcg cgtttctaaa acaaaaacag acaacaacaa cagcaacttt acctggactg 3720 gtgcttcaaa atataacctt aatgggcgtg aatctataat caaccetggc actgctatgg 3780 cctcacacaa agacgacaaa gacaagttct ttcccatgag cggtgtcatg atttttggaa 3840 aggagagege eggagettea aacaetgeat tggacaatgt catgateaea gaegaagagg 3900 aaatcaaagc cactaacccc gtggccaccg aaagatttgg gactgtggca gtcaatctcc 3960 agagcagcag cacagaccct gcgaccggag atgtgcatgt tatgggagcc ttacctggaa 4020 tggtgtggca agacagagac gtatacctgc agggtcctat ttgggccaaa attcctcaca 4080 eggatggaca ettteaceeg teteetetea tgggeggett tggaettaag caceegeete 4140 ctcagatect catcaaaaac acgeetgtte etgegaatee teeggeagag tttteggeta 4200 caaagtttgc ttcattcatc acccagtatt ccacaggaca agtgagcgtg gagattqaat 4260 gggagctgca gaaagaaaac agcaaacgct ggaatcccga agtgcagtat acatctaact 4320 atgcaaaatc tgccaacgtt gatttcactg tggacaacaa tggactttat actgagcctc 4380

<210> 20

<211> 16

<212> DNA

<213> rep binding motif

<400> 20

gctcgctcgc tcgctg

16